

ON PRIMITIVE IDEALS

VICTOR GINZBURG

Abstract

We extend two well-known results on primitive ideals in enveloping algebras of semisimple Lie algebras, the *Irreducibility theorem* for associated varieties and *Duflo theorem* on primitive ideals, to much wider classes of algebras. Our general version of Irreducibility theorem says that if A is a positively filtered associative algebra such that $\text{gr } A$ is a commutative Poisson algebra with *finitely many* symplectic leaves, then the associated variety of any primitive ideal in A is the closure of a single connected symplectic leaf. Our general version of Duflo theorem says that if A is an algebra with a ‘triangular structure’, see §2, then any primitive ideal in A is the annihilator of a simple highest weight module. Applications to Symplectic reflection algebras and Cherednik algebras are discussed.

1 Introduction

Let A be an associative, not necessarily commutative, algebra. Recall that a two-sided ideal $I \subsetneq A$ is called *primitive* if it is the annihilator of a simple left A -module. Primitive ideals often play a role similar to the one played by maximal ideals in the case of commutative algebras. The set of all primitive ideals in A , equipped with Jacobson topology, is called the *primitive spectrum* of A . It is a fundamental invariant of A analogous to the affine scheme $\text{Spec } A$ in the commutative case.

A complete understanding of the primitive spectrum is only available in some ‘very special’ cases. Our motivation in this paper comes from one such case where $A = \mathcal{U}\mathfrak{g}$ is the universal enveloping algebra of a complex semisimple Lie algebra $\mathfrak{g}$. Classification of primitive ideals in $\mathcal{U}\mathfrak{g}$ was one of the central themes of representation theory during the 70–80’s, and it is nowadays well understood, cf. [J1], [J2] and references therein. Yet, details of the classification and the methods involved still appear to be quite complicated.

Our goal is to show that, quite surprisingly, two very important results on primitive ideals in $\mathcal{U}\mathfrak{g}$ have natural generalizations to much wider classes of algebras. The first result is *Duflo theorem* [Du], saying that any primitive ideal in $\mathcal{U}\mathfrak{g}$ is the annihilator of a simple *highest weight* module. We generalize this to arbitrary algebras A that have a triangular structure, i.e. a pair of ‘opposite’ subalgebras $A^\pm$ such that A is a finite type $(A^+ - A^-)$ -bimodule, satisfying certain additional technical conditions, see §2. Our approach is based on the notion of Jacquet functor borrowed from representation theory, and on Gabber’s *Separation theorem* [Ga].

The second result is a generalization of the *Irreducibility theorem* saying that the associated variety of a primitive ideal in $\mathcal{U}\mathfrak{g}$ is irreducible, specifically, it is the closure of a nilpotent conjugacy class in $\mathfrak{g}$. The latter theorem was first partially proved (by a case-by-case argument) in [BoBr], and in a more conceptual way in [KT] and [J2] (independently),

using many earlier deep results due to Joseph, Gabber, Lusztig, Vogan and others. Our argument is an adaptation of a much more direct proof discovered subsequently by Vogan [Vo], combined with some recent results by Brown-Gordon [BrGo].

We got involved into these matters because of our interest in primitive ideals in Symplectic reflection algebras, a new important class of associative algebras introduced recently in [EG] in connection with various works in combinatorics, completely integrable systems, and generalized McKay correspondence for orbifolds $\mathbb{C}^n/\Gamma$. In §6, we will describe all possible subvarieties in $\mathbb{C}^n/\Gamma$ that may arise as associated varieties of primitive ideals in a Symplectic reflection algebra.

An important special class of Symplectic reflection algebras comes from Coxeter groups of finite root systems. The algebras in question have been called *rational Cherednik algebras*, see [EG], [DO], [BEG], since they appeared as a ‘rational’ degeneration of *double-affine Hecke algebras* introduced by Cherednik [Ch]. Thus, the latter may (and should) be thought of as a deformation of the former. From this point of view, representation theory of the rational Cherednik algebra is perhaps ‘more basic’ than (or at least should be studied before) that of the double-affine Hecke algebra in the same sense as the representation theory of semisimple Lie algebras is ‘more basic’ than that of the corresponding quantum groups.

‘Experimental evidence’ suggests that, although rational Cherednik algebras are quite far from enveloping algebras, their representation theory shares many features of representation theory of semisimple Lie algebras, see [BEG]. Thus, classification of primitive ideals in Cherednik algebras seems to be a challenging problem. The results of the present paper provide a first step in this direction.

In more detail, let H_c be the Cherednik algebra associated to the Weyl group W of a root system in a vector space $\mathfrak{h}$. There is an analogue of the Bernstein-Gelfand-Gelfand category of highest weight modules for H_c , to be denoted $\mathcal{O}_c$, see [BEG]. The isomorphism classes of simple objects in $\mathcal{O}_c$ are known to be parametrized by the set $\text{Irrep}(W)$ of irreducible representations of the group W . Given $\sigma \in \text{Irrep}(W)$, let L_σ denote the corresponding simple object of $\mathcal{O}_c$, and $I_\sigma := \text{Ann}(L_\sigma) \subset H_c$ denote its annihilator. The assignment $\Theta : \sigma \mapsto I_\sigma$ gives a map from $\text{Irrep}(W)$ to the set $\text{Prim}(H_c)$ of primitive ideals in H_c , and our general version of Duflo theorem implies that the map Θ is *surjective*, in particular, $\text{Prim}(H_c)$ is a finite set. Moreover, our Irreducibility theorem implies that $\mathcal{V}(I_\sigma)$, the associated variety of I_σ , is the closure of W -saturation of a certain symplectic leaf in $\mathfrak{h} \times \mathfrak{h}^*$. This leads to two important open problems

Problem 1.1 (Classification of primitive ideals). (i) *Describe the fibers of the map $\Theta : \text{Irrep}(W) \longrightarrow \text{Prim}(H_c)$ in a more direct way.*

(ii) *For each $\sigma \in \text{Irrep}(W)$, describe the associated variety of I_σ , in particular, for which pairs $\sigma, \tau \in \text{Irrep}(W)$ does one have: $\mathcal{V}(I_\sigma) = \mathcal{V}(I_\tau)$?*

Remark. The reader should keep in mind that the Cherednik algebra H_c depends on complex parameter(s) ‘ c ’, which is an analogue of ‘central character’ in the Lie algebra case. The solution to Problem 1.1 should depend on the value of ‘ c ’ in an essential way.

As a step towards Problem 1.1(ii), we will compare, in §6, the associated variety of L_σ with that of its annihilator I_σ . Specifically, we show (Theorem 6.8) that the associated variety of any simple highest weight $\mathbf{H}_c$ -module L equals $\mathcal{V}(\text{Ann } L) \cap (\mathfrak{h} \times \{0\})$, the intersection of the associated variety of the annihilator of L (in $\mathbf{H}_c$) with the Lagrangian subspace $\mathfrak{h} \times \{0\} \subset \mathfrak{h} \times \mathfrak{h}^*$.

Acknowledgments. It is a pleasure to acknowledge my debt to Ofer Gabber for many extremely helpful remarks, and for a very careful reading of the manuscript.

2 Main results

Throughout the paper we work with associative algebras over $\mathbb{C}$, the field of complex numbers. We write $\otimes = \otimes_{\mathbb{C}}$.

Let A be a unital $\mathbb{C}$ -algebra equipped with a multiplicative increasing filtration: $0 = A_{-1} \subset A_0 \subset A_1 \subset \dots$, such that $A_i \cdot A_j \subset A_{i+j}$, and $\cup_{j \geq 1} A_j = A$, $1 \in A_0$. Furthermore, we assume that $\text{gr } A = \bigoplus_j A_j/A_{j-1}$, the associated graded algebra, is a finitely generated commutative algebra without zero-divisors, in other words $\text{gr } A \simeq \mathbb{C}[X]$ is the coordinate ring of a (reduced) irreducible affine algebraic variety X . For any (say, left) ideal $I \subset A$, the filtration on A induces one on I , so that $\text{gr } I$ becomes an ideal in $\text{gr } A$. We write $\mathcal{V}(I) \subset X$ for the zero variety of the ideal $\text{gr } I$, usually referred to as the associated variety of I .

Let ℓ be the maximal integer (or ∞ , if A is commutative) such that, for all $i, j \geq 0$ and $a_i \in A_i$, $a_j \in A_j$, one has: $a_i \cdot a_j - a_i \cdot a_j \in A_{i+j-\ell}$.¹ It is well-known that the assignment: $a_i, a_j \mapsto [a_i, a_j] \bmod A_{i+j-\ell-1}$ descends to a canonical Poisson bracket on $\text{gr } A$ that makes X a Poisson algebraic variety. If X is smooth, then one may view X as a complex-analytic manifold equipped with a holomorphic Poisson structure. For each point $x \in X$ one defines $\mathcal{S}_x$, the *symplectic leaf* through x , to be the set of points that could be reached from x by going along Hamiltonian flows.

If X is not necessarily smooth, let $\text{Sing}(X)$ denote the singular locus of X , and for any $k \geq 1$ define inductively: $\text{Sing}^k(X) := \text{Sing}(\text{Sing}^{k-1}(X))$. We get a finite partition $X = \bigsqcup_{k=0}^n X^k$, where the strata $X^k := \text{Sing}^{k-1}(X) \setminus \text{Sing}^k(X)$ are smooth analytic varieties (by definition we put $X^0 = X \setminus \text{Sing}(X)$). It is known, cf. e.g., [BrGo], [Po], that each X^k inherits a Poisson structure, so for any point $x \in X^k$ there is a well-defined symplectic leaf $\mathcal{S}_x \subset X^k$. This way one defines symplectic leaves on an arbitrary Poisson algebraic variety.

In general, each symplectic leaf is a connected smooth analytic (but not necessarily algebraic) subset in X . However, if the algebraic variety X consists of finitely many symplectic leaves only, then it was shown in [BrGo] that each leaf is a smooth irreducible locally-closed algebraic subvariety in X , and partition into symplectic leaves gives an algebraic stratification of X .

Our first main result reads

¹Note that $\ell \geq 1$ since $\text{gr } A$ is commutative.

Theorem 2.1 (Irreducibility theorem). *Assume that the Poisson variety $\text{Spec}(\text{gr } A)$ has only finitely many symplectic leaves. Then, for any primitive ideal $I \subset A$, the variety $\mathcal{V}(I)$ is the closure of a single (connected) symplectic leaf.*

In the classical case of a semisimple Lie algebra $\mathfrak{g}$, given a primitive ideal $I \subset \mathcal{U}\mathfrak{g}$, let $\mathcal{I} := I \cap \mathcal{Z}$ be the intersection of I with $\mathcal{Z}$, the center of $\mathcal{U}\mathfrak{g}$. A standard argument based on a version of Schur lemma, see e.g. [Di], implies that $\mathcal{I}$ is a maximal ideal in $\mathcal{Z}$. We let $A := \mathcal{U}\mathfrak{g}/\mathcal{I} \cdot \mathcal{U}\mathfrak{g}$ be equipped with the filtration induced by the standard increasing filtration on $\mathcal{U}\mathfrak{g}$. The algebraic variety $\text{Spec}(\text{gr } A)$ is known to coincide with the *nilpotent variety* in $\mathfrak{g}$. The latter variety is partitioned into finitely many conjugacy classes. These turn out to be exactly the symplectic leaves. Thus, our Theorem becomes the well-known result first proved in [BoBr, Theorem 6.5] for classical simple Lie algebras, and in [J2, Theorem 3.10] and [KT, Proposition 11] in general (in the Lie algebra case our argument reduces to the proof in [Vo], and doesn't give anything new).

To formulate our second main result, fix a unital associative $\mathbb{C}$ -algebra A , and given $a \in A$ write $\text{ad } a \mapsto [a, x] := a \cdot x - x \cdot a$ for the adjoint action of a on A . Put $\mathbb{Z}_+ = \{1, 2, \dots\}$, and $\mathbb{Z}_- = -\mathbb{Z}_+$.

Now, let $A^\pm \subset A$ be two (non-unital) finitely-generated subalgebras in A and $\delta \in A$, such that

- (T0) A is finitely generated as an $A^+ \otimes (A^-)^{\text{op}}$ -module, i.e., as an $(A^+ \cdot A^-)$ -bimodule.
- (T1) The action of $\text{ad } \delta$ on A is diagonalizable; the subalgebras $A^\pm$ are $\text{ad } \delta$ -stable.
- (T2) All the eigenvalues of $\text{ad } \delta$ -action on $A^\pm$ are in $\mathbb{Z}_\pm$ ('+' for A^+ , and '-' for A^-).

We call the data $(A^\pm, \delta)$ a *commutative triangular structure* on A if conditions (T0)-(T2) hold and, moreover, both subalgebras $A^\pm$ are commutative. We will also consider non-commutative subalgebras $A^\pm$. In that case, we will assume further that the algebra A is equipped with a multiplicative increasing filtration: $0 = A_{-1} \subset A_0 \subset A_1 \subset \dots$. We write $\text{gr } A$ for the associated graded algebra (which is not assumed to be commutative, in general), and endow $A^\pm$ with induced filtrations.

We will say that the data $(A^\pm, \delta)$ gives a *non-commutative triangular structure* on A if, in addition to (T0)-(T2), the following holds:

- (T0') $\text{gr } A$ is a finitely generated $(\text{gr } A^+ \cdot \text{gr } A^-)$ -bimodule.
- (T1') We have: $\delta \in A_1$, moreover, the image of δ in A_1/A_0 is a central element in $\text{gr } A$.
- (T3) The subalgebras $\text{gr } A^\pm$ are commutative.
- (T4) The algebras $\text{gr } A^\pm$ are generated by the subspace $\text{gr}_0 A^\pm \oplus \text{gr}_1 A^\pm$, the corresponding degree ≤ 1 components. Moreover, $\dim(\text{gr}_i A^\pm) < \infty$.

Clearly, (T0') implies (T0). Conditions (T3)-(T4) imply that the subspaces $A_1^\pm$ form finite dimensional Lie subalgebras in A (with respect to the commutator bracket). We denote them by $\mathfrak{n}^\pm = A_1^\pm$. It follows, since $\text{gr } A^\pm$ is generated by the corresponding degree ≤ 1 components, that the algebra $A^\pm$ is generated by the subspace $A_1^\pm$, hence it is the quotient of the augmentation ideal in $\mathcal{U}\mathfrak{n}^\pm$ by some other $\text{ad } \delta$ -stable two-sided ideal in the enveloping algebra $\mathcal{U}\mathfrak{n}^\pm$. Condition (T1') implies that, for each $i \geq 1$, the space

A_i is $\mathbf{ad}\delta$ -stable. Further, the Lie algebras $\mathfrak{n}^\pm$ are necessarily nilpotent, and also $1 \notin A^\pm$ (unless $A = 0$), because of (T2).

Given a non-unital algebra $\mathcal{A}$ and an $\mathcal{A}$ -module M , we say that $\mathcal{A}$ -action on M is locally-nilpotent if, for any $m \in M$, there exists a large enough integer $n = n(m) \gg 0$ such that: $a_1 \cdot \dots \cdot a_n \cdot m = 0$, $\forall a_1, \dots, a_n \in \mathcal{A}$, and similarly for right $\mathcal{A}$ -modules.

Definition 2.2. *Given an algebra A with triangular structure, let ${}^{\uparrow}\mathcal{O}$, resp. $\mathcal{O}_{\downarrow}$, be the category of finitely-generated left, resp. right, A -modules M such that the A^- -action, resp. A^+ -action, on M is locally-nilpotent.*

Recall that a two-sided ideal $I \subsetneq A$ is called prime, if for any two-sided ideals $J_1, J_2 \subset A$ we have: $J_1 \cdot J_2 \subset I \implies J_1 \subset I$ or $J_2 \subset I$. Our second main result reads

Theorem 2.3 (Generalized Duflo theorem). *Let $(A^\pm, \delta)$ be a (commutative or non-commutative) triangular structure on A . Assume that the category ${}^{\uparrow}\mathcal{O}$ (or $\mathcal{O}_{\downarrow}$) has finitely many isomorphism classes of simple objects. Then the following conditions on a two-sided ideal $I \subsetneq A$ are equivalent:*

- (i) *I is a primitive ideal;*
- (ii) *I is a prime ideal;*
- (iii) *I is the left annihilator of a simple object of ${}^{\uparrow}\mathcal{O}$.*
- (iv) *I is the right annihilator of a simple object of $\mathcal{O}_{\downarrow}$.*

In the classical case, one considers a semisimple Lie algebra $\mathfrak{g}$ with a fixed triangular decomposition: $\mathfrak{g} = \mathfrak{n}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}^-$. Given a primitive ideal $I \subset \mathcal{U}\mathfrak{g}$, we define $A := \mathcal{U}\mathfrak{g}/\mathcal{I} \cdot \mathcal{U}\mathfrak{g}$, where $\mathcal{I} := I \cap \mathcal{Z}$ as above. Let $A^\pm$ be the image in A of $\mathfrak{n}^\pm \cdot \mathcal{U}\mathfrak{n}^\pm$. The Harish-Chandra homomorphism: $\mathcal{Z} \hookrightarrow \mathrm{Sym} \mathfrak{h}$, combined with Poincaré-Birkhoff-Witt theorem imply easily that A is a finitely generated $(A^+ \cdot A^-)$ -bimodule, see e.g. [Di]. We equip A with an increasing filtration induced by the standard increasing filtration on $\mathcal{U}\mathfrak{g}$. Further, if $\delta \in \mathfrak{h}$ denotes the sum of positive coroots, then the $\mathbf{ad}\delta$ -action on $\mathcal{U}\mathfrak{n}^\pm$ is known to be diagonal with eigenvalues in $\mathbb{Z}_\pm$. The data $(A^\pm, \delta)$ gives a non-commutative triangular structure on A . The categories ${}^{\uparrow}\mathcal{O}$ and $\mathcal{O}_{\downarrow}$ are both equivalent, in this case, to the standard category $\mathcal{O}$ (with fixed central character corresponding to $\mathcal{I}$) as defined by Bernstein-Gelfand-Gelfand. Simple objects of the latter category are well-known to be parametrized by a (necessarily finite) Weyl group orbit, so that Theorem 2.3 applies. This gives a new proof (cf. also [J2, n.2.4] for a similar idea) of the original Duflo theorem [Du].

Remark. In the special case $A = \mathcal{U}\mathfrak{g}$ above, our proof of Duflo theorem is perhaps not much simpler than the earlier proofs, see [Du], [BG], [J1], because of its heavy dependence on the difficult Separation theorem due to Gabber [Ga]. One advantage of our approach however is that, unlike all other proofs, it requires very little information about representation theory of the algebra A .

Simultaneously with Theorem 2.3 we will also prove

Theorem 2.4. *Under the assumptions of Theorem 2.3, the algebra A has finite length as an $A \otimes A^{\mathrm{op}}$ -module.*

We say that a δ -action on a vector space M is locally-finite, if any element $m \in M$ is contained in a δ -stable finite dimensional vector subspace.

Our last important result reads

Theorem 2.5. *Let $(A^\pm, \delta)$ be a triangular structure on A . Then*

(i) *The δ -action on any object M of either ${}^1\mathcal{O}$ or $\mathcal{O}_\downarrow$ is locally-finite, and all (generalized) eigenspaces of this δ -action are finite-dimensional.*

(ii) *The following two properties of the category ${}^1\mathcal{O}$, resp. $\mathcal{O}_\downarrow$, are equivalent*

- *The category has only finitely many (isomorphism classes of) simple objects.*
 - *Any object of the category has finite length.*
- (2.6)

(iii) *If properties (2.6) hold for ${}^1\mathcal{O}$ or $\mathcal{O}_\downarrow$, then both categories ${}^1\mathcal{O}$, $\mathcal{O}_\downarrow$, have enough projectives and injectives.*

3 Proof of Theorem 2.1

We need to recall some results due to Brown-Gordon [BrGo].

Let B be a finitely generated Poisson algebra, with Poisson bracket $\{-, -\}$. An ideal $J \subset B$ (with respect to the commutative product) is called a *Poisson ideal* if it is also an ideal with respect to the Lie bracket $\{-, -\}$, i.e. if $\{J, B\} \subset J$. For the following standard result of commutative algebra, see e.g. [CO, Theorem 4.5], or [Di, §3.3.2].

Lemma 3.1. *If J is a Poisson ideal then so are both its radical and all the associated prime ideals in B .* □

Write $\mathfrak{m}_x$ for the maximal ideal in B corresponding to a closed point $x \in \operatorname{Spec} B$, and recall the notation $\mathcal{S}_x \subset \operatorname{Spec} B$ for the symplectic leaf through x . Following [BrGo], given any ideal $J \subset B$ we let $\mathcal{P}(J)$ denote the maximal Poisson ideal contained in J . If J is prime then so is $\mathcal{P}(J)$, and one has

Proposition 3.2 ([BrGo], Prop. 3.7). *Assume the variety $\operatorname{Spec} B$ consists of finitely many symplectic leaves. Then, for any closed point $x \in \operatorname{Spec} B$, the leaf $\mathcal{S}_x$ coincides with the regular locus of the zero variety of the ideal $\mathcal{P}(\mathfrak{m}_x)$. Thus, $\mathcal{S}_x$ is a smooth connected locally-closed subvariety in $\operatorname{Spec} B$.* □

Recall that the Poisson structure on B restricts to a (non-degenerate) symplectic structure on each symplectic leaf. In particular, all symplectic leaves have even (complex) dimension. Hence, from Lemma 3.1 and Proposition 3.2 we deduce

Corollary 3.3. *If the variety $\operatorname{Spec} B$ consists of finitely many symplectic leaves, then every irreducible component of the zero variety of any Poisson ideal $J \subset B$ is the closure of a symplectic leaf, in particular, it is even-dimensional.* □

Next, let A be an associative filtered $\mathbb{C}$ -algebra as in Theorem 2.1. We recall

Lemma 3.4 ([BoKr], Korollar 3.6). *If $I \subset J \subset A$ are two-sided ideals, $I \neq J$ and moreover I is prime, then: $\dim \mathcal{V}(J) < \dim \mathcal{V}(I)$. $\square$*

Given a finitely generated A -module M , choose a *good* filtration (cf. e.g. [Be]) on M and write $\text{Supp}(\text{gr } M)$ for the support of the corresponding associated graded $\text{gr } A$ -module, a reduced algebraic subvariety in $\text{Spec}(\text{gr } A)$. It is well-known (due to Bernstein [Be]) that this subvariety is independent of the choice of a good filtration on M , and it will be denoted $\text{Supp } M$ below. Note that if $I \subset A$ is a left ideal, then in the special case: $M = A/I$ we have by definition $\text{Supp}(A/I) = \mathcal{V}(I)$.

Assume from now on that $\text{gr } A$ is a Poisson algebra with finitely many symplectic leaves. It is straightforward to verify that, for any two-sided ideal $I \subset A$, the associated graded ideal, $\text{gr } I$, is a Poisson ideal in $\text{gr } A$. Thus by Corollary 3.3, every irreducible component of $\mathcal{V}(I)$, the zero variety of $\text{gr } I$, is the closure of a symplectic leaf.

Proof of Theorem 2.1: Let I be a primitive ideal. We've just seen that proving the Theorem amounts to showing that $\mathcal{V}(I)$ is irreducible. To prove this, we mimic the argument in ([Vo], §§3-4). Put $M = A/I$, viewed as an A -bimodule, and let $X := \text{Spec}(\text{gr } A)$. The filtration on A induces a filtration on M , and we view $\text{gr } M = \text{gr } A / \text{gr } I$ as a finitely generated $\text{gr } A$ -module. Let $\mathcal{M}$ denote the coherent sheaf on X corresponding to this $\text{gr } A$ -module. By definition we have: $\text{Supp } \mathcal{M} = \mathcal{V}(I)$, and we must show that this variety is irreducible. Pick an irreducible component of $\text{Supp } \mathcal{M}$ of maximal dimension, $\dim(\text{Supp } \mathcal{M})$. Corollary 3.3 says that there exists a symplectic leaf $\mathcal{S} \subset X$ such that this irreducible component is $\overline{\mathcal{S}}$, the Zariski closure of $\mathcal{S}$. Furthermore, by Lemma 3.1, any imbedded component² of $\mathcal{V}(I)$ (corresponding to an associated prime of $\text{gr } I$) that has non-empty intersection with $\mathcal{S}$ must contain the whole of $\mathcal{S}$.

A key point is that, since all symplectic leaves are even dimensional, we have

$$\dim(\overline{\mathcal{S}} \setminus \mathcal{S}) \leq \dim \mathcal{S} - 2 = \dim(\text{Supp } \mathcal{M}) - 2. \quad (3.5)$$

The idea of the proof of Irreducibility Theorem given in ([Vo], §4), that we will follow closely, is to introduce a certain new filtration, $F_\bullet$, on $M = A/I$ defined "locally" in terms of the stratum $\mathcal{S}$, and such that $\text{gr }^F M$ gives rise to a coherent sheaf on $\mathcal{S}$. Then, since the boundary of $\mathcal{S}$ has codimension two by (3.5), the global sections of $\text{gr }^F M$ over $\mathcal{S}$ necessarily form a *finitely generated* $\text{gr } A$ -module which is, moreover, automatically supported on $\overline{\mathcal{S}}$. Hence, the filtration F is good, and one obtains $\mathcal{V}(I) = \text{Supp}(A/I) = \text{Supp}(\text{gr }^F A/I) = \overline{\mathcal{S}}$.

We proceed to a more detailed exposition. First, Proposition 3.2 implies that $\mathcal{S}$ is Zariski-open in $\overline{\mathcal{S}}$, hence also in $\text{Supp } \mathcal{M}$. Therefore, $Z := (\text{Supp } \mathcal{M}) \setminus \mathcal{S}$ is a Zariski closed subset in X . Thus $X \setminus Z$ is a Zariski open subset in X . We may choose a smaller *affine* Zariski open subset $U \subset X \setminus Z$ such that $(\text{Supp } \mathcal{M}) \cap U = \mathcal{S}$. Write $j : U \hookrightarrow X$ for the corresponding open imbedding. Set $\mathcal{M}_U := \mathcal{M}|_U$, a coherent sheaf on U . By construction, the leaf $\mathcal{S}$ is closed in U , and $\text{Supp } \mathcal{M}_U = \mathcal{S}$. Hence, $\text{Supp}(j_* \mathcal{M}_U) = \overline{\mathcal{S}}$. Therefore, (3.5) yields

$$\dim(\text{Supp}(j_* \mathcal{M}_U) \cap (X \setminus U)) \leq \dim(\text{Supp } \mathcal{M}_U) - 2.$$

² here $\mathcal{V}(I)$ is viewed as a scheme rather than as a reduced variety.

A standard result of algebraic geometry now says that $j_*\mathcal{M}_U$ is a coherent sheaf on X . Therefore, we have:

$$\Gamma(U, \mathcal{M}_U) = \Gamma(X, j_*\mathcal{M}_U) \text{ is a finitely generated } \mathbf{gr} A\text{-module.} \quad (3.6)$$

We are going to apply microlocal techniques – pioneered by Gabber [Ga1] – equivalent to the argument in ([Vo], §3), that has been also inspired by Gabber. To this end, observe that the variety X has a cone-structure, that is a contracting $\mathbb{C}^*$ -action induced by the (non-negative) grading on $\mathbf{gr} A$. Clearly, $\mathcal{V}(I)$, hence $\mathcal{S}$, Z , are $\mathbb{C}^*$ -stable subvarieties in X .

For any non-zero homogeneous element $f \in \mathbf{gr} A$, let $\mathcal{U}_f$ denote the complement in X of the divisor $f = 0$. Thus, $\mathcal{U}_f$ is an affine Zariski open $\mathbb{C}^*$ -stable subset of X . Let $\widehat{A}_{\mathcal{U}_f}$ be the formal micro-localization of A at $\mathcal{U}_f$, see e.g. ([Gi], §1.3). Thus, $\widehat{A}_{\mathcal{U}_f}$ is a complete $\mathbb{Z}$ -filtered ring such that $\mathbf{gr}(\widehat{A}_{\mathcal{U}_f}) = \mathbb{C}[\mathcal{U}_f]$, the ring of regular functions on the affine open set $\mathcal{U}_f$. The assignment: $\mathcal{U}_f \mapsto \widehat{A}_{\mathcal{U}_f}$ extends to a sheaf $\mathcal{U} \mapsto \widehat{A}_{\mathcal{U}}$ in the topology of Zariski open cone-subsets in $X \setminus \{0\}$. For any Zariski open cone-subset $\mathcal{U}$, we have an imbedding $\mathbf{gr}(\widehat{A}_{\mathcal{U}}) \hookrightarrow \mathbb{C}[\mathcal{U}]$, which becomes an equality if $\mathcal{U}$ is affine. There is a canonical algebra imbedding $A \hookrightarrow \widehat{A}_{\mathcal{U}}$, which is strictly compatible with the filtrations and such that the induced map $\mathbf{gr} A \rightarrow \mathbf{gr}(\widehat{A}_{\mathcal{U}})$ gets identified with restriction of functions: $\mathbf{gr} A = \mathbb{C}[X] \rightarrow \mathbb{C}[\mathcal{U}]$.

Further, given a finitely generated A -module L , we let $\widehat{L}_{\mathcal{U}_f} = \widehat{A}_{\mathcal{U}_f} \otimes_A L$ denote its formal micro-localization, a finitely generated $\widehat{A}_{\mathcal{U}_f}$ -module. The assignment: $\mathcal{U}_f \mapsto \widehat{L}_{\mathcal{U}_f}$ extends to a sheaf $\widehat{L}$ in the topology of Zariski open cone-subsets in $X \setminus \{0\}$. One has a canonical sheaf isomorphism $\widehat{L} \simeq \widehat{A} \otimes_A L$, in particular, an isomorphism $\Gamma(\mathcal{U}, \widehat{L}) \simeq \Gamma(\mathcal{U}, \widehat{A}) \otimes_A L$, for any *affine* open cone-subset $\mathcal{U}$.

We now set $\mathcal{U} := U$, the affine Zariski open subset in $X \setminus Z$ that has been chosen earlier, cf. (3.6), and which we now additionally assume to be a cone-subset. We apply the formal micro-localization at U to our A -module $M = A/I$, to get the left $\widehat{A}_U$ -module $\widehat{M}_U = \widehat{A}_U \otimes_A (A/I) = \widehat{A}_U / \widehat{A}_U \cdot I$. We note that since M has also a right A -action (the ideal I is a two-sided ideal) the left $\widehat{A}_U$ -module $\widehat{M}_U$ acquires a canonical $\widehat{A}_U$ -bimodule structure. Equivalently, one can get the same bimodule by applying the formal micro-localization procedure to M viewed as a left $A \otimes A^{\text{op}}$ -module (from this point of view, $\widehat{M}_U$ is ‘supported’ on the diagonal $U \subset U \times U$, since M , viewed as a $A \otimes A^{\text{op}}$ -module, is clearly ‘supported’ on the diagonal $X \subset X \times X$). Further, we have a canonical A -bimodule map

$$i: M \rightarrow \widehat{A}_U \otimes_A M = \widehat{M}_U.$$

Let K denote the kernel of i and $\overline{M}$ denote the image of i . Both K and $\overline{M}$ are A -bimodules again, in particular, the preimage of K under the projection $A \twoheadrightarrow A/I = M$ is clearly a two-sided ideal, say $J \subset A$. Thus, we have the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & J/I & \longrightarrow & A/I & \longrightarrow & A/J \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ & & K & \longrightarrow & M & \longrightarrow & \overline{M} \end{array} \xrightarrow{i} \widehat{M}_U = \widehat{A}_U \otimes_A M, \quad (3.7)$$

where we have factored the map i as a composition: $M \twoheadrightarrow \overline{M} \xrightarrow{\iota} \widehat{M}_U$. The latter factorization shows that the imbedding ι induces an isomorphism: $\widehat{A}_U \otimes_A \overline{M} \xrightarrow{\sim} \widehat{A}_U \otimes_A M$ of the corresponding micro-localizations. The micro-localization functor being exact, this yields $\widehat{K}_U = \widehat{A}_U \otimes_A K = 0$. Hence $(\mathcal{S}upp K) \cap U = \emptyset$.

We claim that $K = 0$. If not, then the two-sided ideal $J \subset A$, see (3.7), properly contains I . Hence, Lemma 3.4 yields $\dim(\mathcal{S}upp A/J) < \dim(\mathcal{S}upp A/I)$. But since $\overline{\mathcal{S}}$ is a component of $\mathcal{S}upp A/I$ of maximal dimension, we get: $\dim \overline{\mathcal{S}} > \dim(\mathcal{S}upp A/J)$. Therefore, the equality of sets: $\mathcal{S}upp A/I = (\mathcal{S}upp J/I) \cup (\mathcal{S}upp A/J)$, which follows from the short exact sequence in the top row of (3.7), cannot be achieved unless $\mathcal{S} \subset \mathcal{S}upp J/I = \mathcal{S}upp K$. This last inclusion contradicts the conclusion of the previous paragraph, saying that $(\mathcal{S}upp K) \cap U = \emptyset$. Thus, $K = 0$ and our claim follows.

To proceed further, observe that the module $\widehat{M}_U$ comes equipped with a natural increasing filtration $F_\bullet \widehat{M}_U$ obtained, essentially, by localizing the standard filtration on $M = A/I$. Moreover, for the associated graded module corresponding to the filtration $F_\bullet \widehat{M}_U$ one has a canonical isomorphism $\mathbf{gr}^F(\widehat{M}_U) \xrightarrow{\sim} \Gamma(U, \mathcal{M}_U)$, since U is affine. We see that $\mathbf{gr}^F(\widehat{M}_U)$ is a finitely generated $\mathbf{gr} A$ -module, due to (3.6). Further, the increasing filtration on $\widehat{M}_U$ induces a filtration $F_\bullet M = M \cap F_\bullet \widehat{M}_U$ via the imbedding $i: M \hookrightarrow \widehat{M}_U$ (we have shown that i is injective). Then $\mathbf{gr}^F M$ is a $\mathbf{gr} A$ -submodule of $\mathbf{gr}^F(\widehat{M}_U)$, hence is also finitely generated. Therefore the filtration $F_\bullet M$ is a *good* filtration on M . This yields $\mathcal{S}upp M = \mathcal{S}upp(\mathbf{gr}^F M)$, and we deduce

$$\mathcal{S} \subset \mathcal{V}(I) = \mathcal{S}upp M = \mathcal{S}upp(\mathbf{gr}^F M) \subset \mathcal{S}upp(\mathbf{gr}^F \widehat{M}_U) \subset \mathcal{S}upp(j_* \mathcal{M}_U) \subset \overline{\mathcal{S}}.$$

It follows that $\mathcal{V}(I) = \overline{\mathcal{S}}$, and the Theorem is proved. $\square$

4 Basic properties of categories $\mathcal{O}^\uparrow$ and $\mathcal{O}_\downarrow$

In this section we prove a few technical lemmas that will be of use later.

Given a δ -action on a vector space V and $\mu \in \mathbb{C}$, for the generalized μ -eigenspace of δ write $V_\mu := \{v \in V \mid (\delta - \mu \cdot \text{Id})^N \cdot v = 0 \text{ for } N \gg 0\}$. If the action on V is locally-finite then there is a (possibly infinite) direct sum decomposition $V = \bigoplus_{\mu \in \mathbb{C}} V_\mu$.

Let $(A^\pm, \delta)$ be a triangular structure on A . We consider the following statements about an A -module M :

1. M is finitely generated over A^+ . Furthermore, for any good filtration on M (4.1) as an A -module, $\mathbf{gr} M$ is finitely-generated over $\mathbf{gr}(A^+)$.
2. The δ -action on M is locally-finite and each generalized δ -eigenspace M_μ is finite-dimensional.
3. The real parts of the eigenvalues of δ -action in M are all bounded from below.

It will be convenient below to introduce the notation ${}^\heartsuit A^\pm := A^\pm + \mathbb{C} \cdot 1 \subseteq A$ and, for $m \geq 1$, put $(A^-)^m := A^- \cdot \dots \cdot A^-$ (m factors).

Lemma 4.2. *Let M be an A -module generated by a finite-dimensional subspace $M_0 \subset M$ such that, for some $m \geq 1$, we have: $(A^-)^m \cdot M_0 = 0$. Then properties (4.1)(1)-(3) hold.*

Proof. We start with statement (4.1)(2). Properties (T0)-(T1) imply that there exists an $\mathbf{ad}\delta$ -stable finite-dimensional $\mathbb{C}$ -vector subspace $E \subset A$ such that $A = {}^\heartsuit A^+ \cdot E \cdot {}^\heartsuit A^-$. Furthermore, looking at the weights of $\mathbf{ad}\delta$ -action on A , for the centralizer of δ in A we find:

$$A^{\mathbf{ad}\delta} \quad (:= \{a \in A \mid [a, \delta] = 0\}) = \sum_{\beta, \gamma \in \mathbb{C}} ({}^\heartsuit A^+)_{-\beta-\gamma} \cdot E_\beta \cdot ({}^\heartsuit A^-)_\gamma, \quad (4.3)$$

where $(-)_\beta$ denotes $\mathbf{ad}\delta$ -eigenspace in $(-)$ with the eigenvalue β .

Observe further that, for any integer $m > 0$, the space $(A^-)^m$ has finite codimension in A^- since a similar result holds for the enveloping algebra of a (finite-dimensional) nilpotent Lie algebra. It follows that, for all $\gamma \in \mathbb{C}$ except possibly a finite set, one has: $({}^\heartsuit A^-)_\gamma \subset (A^-)^m$. Further, the algebra A^+ being finitely-generated, each eigenspace $(A^+)_\alpha$ is finite-dimensional, by (T3). Therefore, (4.3) implies that there exist finite-dimensional subspaces $B_m \subset A^\delta$ such that

$$A^{\mathbf{ad}\delta} = B_m + A^{\mathbf{ad}\delta} \cap ({}^\heartsuit A^+ \cdot E \cdot (A^-)^m) \quad , \quad m = 1, 2, \dots$$

Hence, since $(A^-)^m \cdot M_0 = 0$, we get $A^{\mathbf{ad}\delta} \cdot M_0 = B_m \cdot M_0$, is a finite dimensional space. The subspace $A^{\mathbf{ad}\delta} \cdot M_0 \subset M$ being clearly δ -stable, we conclude that M is generated as an A -module by the δ -stable finite dimensional subspace $A^{\mathbf{ad}\delta} \cdot M_0 \subset M$. From $M = A \cdot M_0 \subset A \cdot (A^{\mathbf{ad}\delta} \cdot M_0)$ we deduce that the δ -action on M is locally-finite, since $\mathbf{ad}\delta$ -action on A is locally-finite.

To prove statement (4.1)(3), choose B , a complementary subspace to $(A^-)^m$ in ${}^\heartsuit A^-$ so that ${}^\heartsuit A^- = B \oplus (A^-)^m$. Then we have: $M = A \cdot M_0 = {}^\heartsuit A^+ \cdot E \cdot (B \oplus (A^-)^m) \cdot M_0 = {}^\heartsuit A^+ \cdot E \cdot B \cdot M_0$. Further, the space $E \cdot B \cdot M_0$ is finite-dimensional, and the action of A^+ may only increase the real parts of the eigenvalues of δ -action in M . It follows that the real parts of the eigenvalues of δ -action in M are all bounded from below, and (4.1)(3) is proved.

Next, since each $\mathbf{ad}\delta$ -eigenspace in A^+ is finite-dimensional, we deduce from the equality $M = {}^\heartsuit A^+ \cdot (E \cdot B \cdot M_0)$ that each δ -eigenspace in M is finite-dimensional. This completes the proof of (4.1)(2). The same equality clearly implies that M is finitely generated over A^+ .

To complete the proof of (4.1)(1) we recall, see e.g. [CG, Corollary 2.3.19], the following well-known fact: if for *some* good filtration on M , the associated graded module $\mathbf{gr} M$ is finitely-generated over $\mathbf{gr} A^+$, then the same holds for *any* good filtration on M . Thus, we may prove the result for a particular good filtration on M of our choice. To this end, observe that M is, by definition, a quotient of the direct sum of finitely many copies of the A -module $A/A \cdot (A^-)^m$. We equip $A/A \cdot (A^-)^m$ with the filtration inherited from the standard filtration on A . Further, equip the direct sum with the direct sum filtration, and choose the filtration on M to be the induced quotient-filtration. This way, we see that $\mathbf{gr} M$ is a quotient of the direct sum of finitely many copies of the $\mathbf{gr} A$ -module

$(\text{gr } A)/(\text{gr } A) \cdot \text{gr } (A^-)^m$. Thus, we are reduced to showing that $(\text{gr } A)/(\text{gr } A) \cdot (\text{gr } A^-)^m$ is a finitely generated $\text{gr } A^+$ -module. But this is clear from property (T0'), saying that $\text{gr } A$ is a finitely generated $(\text{gr } A^+)$ - $(\text{gr } A^-)$ -bimodule. $\square$

Recall the categories ${}^1\mathcal{O}$ and $\mathcal{O}_\downarrow$ from Definition 2.2.

Corollary 4.4. *An A -module M is generated by a finite-dimensional subspace $M_0 \subset M$ such that $(A^-)^m \cdot M_0 = 0$, for some $m \geq 1$, if and only if $M \in {}^1\mathcal{O}$.*

Proof. Assume that M is generated by M_0 , as above. Then Lemma 4.2 implies that statement (4.1)(3) holds. Hence, since the A^- -action decreases the real parts of the eigenvalues of δ -action in M , we conclude that the A^- -action on M is locally-nilpotent.

Conversely, any object of $M \in {}^1\mathcal{O}$ is generated by a finite-dimensional subspace $M_0 \subset M$, and the action of A^- on M is locally-nilpotent. Hence, there exists $m \gg 0$ such that $(A^-)^m \cdot M_0 = 0$. $\square$

Corollary 4.5. *The category ${}^1\mathcal{O}$ is abelian, and properties (4.1)(1)-(3) hold for any object $M \in {}^1\mathcal{O}$.*

Proof. The last claim is immediate from Lemma 4.2 and Corollary 4.4. In particular, we conclude that any object $M \in {}^1\mathcal{O}$ is finitely generated over A^+ , due to property (4.1)(1). Now, A^+ , being a quotient of $\mathfrak{n}^+ \cdot \mathcal{U}\mathfrak{n}^+$, is a Noetherian algebra. Therefore, any A -submodule of M belongs to ${}^1\mathcal{O}$. Hence, the category ${}^1\mathcal{O}$ is abelian. $\square$

Lemma 4.6. *The A -module $A/A \cdot \mathfrak{n}^-$ is an object of ${}^1\mathcal{O}$, and any simple object of ${}^1\mathcal{O}$ is isomorphic to a quotient of $A/A \cdot \mathfrak{n}^-$.*

Proof. The module $A/A \cdot \mathfrak{n}^-$ is generated over A by a single vector (= coset of $1 \in A$), which is annihilated by the algebra A^- . Hence, Corollary 4.4 yields $A/A \cdot \mathfrak{n}^- \in {}^1\mathcal{O}$.

Now, given a simple object $L \in {}^1\mathcal{O}$, we may choose a non-zero element $v \in L$ in the generalized δ -eigenspace corresponding to an eigenvalue with minimal possible real part. Commutation relations show that the action of any element of $\mathfrak{n}^-$ decreases the real part of the eigenvalue. Hence, we must have: $\mathfrak{n}^- \cdot v = 0$. Therefore, the assignment: $a \mapsto a(v)$ gives a well-defined A -module map $A/A \cdot \mathfrak{n}^- \rightarrow L$. This map is nonzero, hence, surjective since L is simple. $\square$

Remark. Analogues of the above results hold also for the category $\mathcal{O}_\downarrow$. $\diamond$

Given a (possibly infinite dimensional) vector space V with locally finite δ -action such that all δ -eigenspaces V_λ are finite-dimensional, let $V^* = \bigoplus_\lambda (V_\lambda)^* \subset \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$ denote the direct sum of the spaces dual to V_λ 's. If V is a left A -module, then V^* is an A -stable subspace in the right A -module $\text{Hom}_{\mathbb{C}}(V, \mathbb{C})$ (= linear dual of V). The right A -module V^* is called the *restricted dual* of V . There is a canonical A -module isomorphism: $V \xrightarrow{\sim} (V^*)^*$.

Lemma 4.7. *Assume that any object of either ${}^1\mathcal{O}$ or $\mathcal{O}_\downarrow$ has finite length. Then the restricted duality functor $M \mapsto M^*$ induces anti-equivalences: ${}^1\mathcal{O} \xleftarrow{\sim} \mathcal{O}_\downarrow$. $\square$*

Proof. An object has finite length if and only if it satisfies both ascending and descending chain conditions. Restricted duality clearly interchanges these two conditions. Thus, $M^\star$ has finite length if and only if so does M .

Further, it is clear that the real parts of the eigenvalues of δ -action in M are bounded from below if and only if the real parts of the eigenvalues of δ -action in $M^\star$ are bounded from above. Thus, it follows by Corollary 4.4 and Lemma 4.2 that $M \in {}^\uparrow\mathcal{O}$ if and only if $M^\star \in \mathcal{O}_\downarrow$. $\square$

Corollary 4.8. *The equivalent conditions (2.6) hold for the category ${}^\uparrow\mathcal{O}$ if and only if they hold for the category $\mathcal{O}_\downarrow$.* $\square$

Proof of Theorem 2.5. Part (i) of the Theorem is immediate from Corollary 4.4 and Lemma 4.2. To prove part (ii), fix $M \in {}^\uparrow\mathcal{O}$ and a simple object $L \in {}^\uparrow\mathcal{O}$. We claim that there exists an integer $a(M : L)$ such that the following holds:

For any descending, resp. ascending, chain of subobjects: $M = M^0 \supset M^1 \supset \dots \supset M^N$, let I be the set formed by the indices ‘ i ’ such that M^i/M^{i+1} contains L as a subquotient. Then, one has: $\#I \leq a(M : L)$. (4.9)

To prove the claim, choose $\mu \in \mathbb{C}$ such that the eigenspace L_μ is non-zero. Then, we clearly have

$$\dim M_\mu = \sum_{i=1}^N \dim(M^i/M^{i+1})_\mu \leq \sum_{i \in I} \dim(M^i/M^{i+1})_\mu \leq \#I \cdot \dim L_\mu.$$

Since $\dim L_\mu \neq 0$, and $\dim M_\mu < \infty$ by part (i) of the Proposition, we deduce: $\#I \leq \dim M_\mu / \dim L_\mu$. Thus, we can take $a(M : L)$ to be the integral part of $(1 + \dim M_\mu / \dim L_\mu)$.

Now, assume that $\{L_1, \dots, L_d\}$ is a complete (finite) collection of isomorphism classes of simple objects of ${}^\uparrow\mathcal{O}$. To prove that an object $M \in {}^\uparrow\mathcal{O}$ has finite length, one has to verify both ascending and descending chain conditions for M . Consider, for instance, a descending chain of subobjects: $M = M^0 \supsetneq M^1 \supsetneq \dots \supsetneq M^N$. We must show that the length of this chain is bounded from above by a certain number $a = a(M)$ which is independent of the chain. But this is clear from Claim 4.9, since for each $i = 1, \dots, N$, the non-zero object M^i/M^{i+1} must contain at least one of the simple objects L_r , $r = 1, \dots, d$, as a subquotient. Therefore, $N \leq \sum_{r=1}^d a(M : L_r)$, and we may put $a(M) := \sum_{r=1}^d a(M : L_r)$.

To complete the proof of part (ii) of the Theorem, assume any object of ${}^\uparrow\mathcal{O}$ has finite length. Then, the module $A/A \cdot \mathfrak{n}^-$ has finite length by Lemma 4.6, hence it follows from the Lemma that the number of isomorphism classes of simple objects in ${}^\uparrow\mathcal{O}$ is finite.

To prove part (iii), write again $\{L_1, \dots, L_d\}$ for a complete collection of isomorphism classes of simple objects of ${}^\uparrow\mathcal{O}$. We know that for each $r = 1, \dots, d$ there is a real number $\lambda_r \in \mathbb{R}$ such that the real parts of δ -eigenvalues in L_r are all $\geq \lambda_r$. It follows that, for any object $M \in {}^\uparrow\mathcal{O}$, the real parts of δ -eigenvalues in M are all $\geq \min(\lambda_1, \dots, \lambda_d)$. Thus, since A^- -action decreases the real parts of δ -eigenvalues by at least 1, we observe that, for each $\mu \in \mathbb{C}$ there exists an integer $n(\mu) > 0$ such that we have:

$$(A^-)^{n(\mu)} \cdot M_\mu = 0 \quad , \quad \forall M \in {}^\uparrow\mathcal{O}. \quad (4.10)$$

Given $\mu \in \mathbb{C}$, for each $k = 1, 2, \dots$, we put

$$J(\mu, k) := A \cdot (\delta - \mu)^k + A \cdot (A^-)^{n(\mu)}.$$

Clearly $J(\mu, 1) \supseteq J(\mu, 2) \supseteq \dots$, is a descending chain of left ideals in A . We claim that this chain stabilizes, i.e., there exists $\ell \gg 0$ such that we have: $J(\mu, \ell) = J(\mu, \ell + 1) = \dots$. To prove the claim, observe first that the left A -module $A/A \cdot (A^-)^{n(\mu)}$ is an object of ${}^1\mathcal{O}$, by Corollary 4.4. Hence, this object has finite length, by part (ii) of the Theorem. Now, for $k = 1, 2, \dots$, let $N^k \subset A/A \cdot (A^-)^{n(\mu)}$ be an A -submodule generated by the coset $(\delta - \mu)^k \bmod (A \cdot (A^-)^{n(\mu)})$. Clearly $N^1 \supseteq N^2 \supseteq \dots$. We conclude that this chain of submodules must stabilize, hence there exists $\ell \gg 0$ such that the canonical projections $(A/A \cdot (A^-)^{n(\mu)})/N^\ell \leftarrow (A/A \cdot (A^-)^{n(\mu)})/N^{\ell+1} \leftarrow \dots$ are all bijections. But these projections are nothing but the canonical projections $A/J(\mu, \ell) \leftarrow A/J(\mu, \ell + 1) \leftarrow \dots$, and our stabilization claim follows.

We put $P(\mu) := A/J(\mu, \ell)$. The A -module $P(\mu)$ is a quotient of $A/A \cdot (A^-)^{n(\mu)}$, hence, an object of ${}^1\mathcal{O}$. We claim³ that $P(\mu)$ is a projective in ${}^1\mathcal{O}$. To see this, let $M \in {}^1\mathcal{O}$ and fix $v \in M_\mu$. Then $(A^-)^{n(\mu)} \cdot v = 0$, due to (4.10). Furthermore, there exists k large enough, such that $(\delta - \mu)^k \cdot v = 0$. Therefore, $J(\mu, k) \cdot v = 0$, hence $J(\mu, \ell) \cdot v = 0$, by the stabilization claim proved above. Therefore, the assignment $a \mapsto a(v)$ descends to a well-defined A -module map $\varphi_v : P(\mu) = A/J(\mu, \ell) \rightarrow M$. Thus, we get a canonical isomorphism

$$M_\mu \xrightarrow{\sim} \text{Hom}_A(P(\mu), M) \quad , \quad v \mapsto \varphi_v. \quad (4.11)$$

We conclude that the functor: ${}^1\mathcal{O} \rightarrow \text{Vector spaces}$, $M \mapsto M_\mu$, which is clearly exact, is represented by the object $P(\mu)$. Hence, $P(\mu)$ is projective.

To complete the proof, let $L \in {}^1\mathcal{O}$ be a simple object. Find $\mu \in \mathbb{C}$ such that $L_\mu \neq 0$. Then (4.11) implies that there is a nonzero A -module map $P(\mu) \rightarrow L$. This map is surjective, since L is simple. Thus, we have shown that any simple object in ${}^1\mathcal{O}$ is isomorphic to a quotient of $P(\mu)$, for an appropriate $\mu \in \mathbb{C}$. It follows that the category ${}^1\mathcal{O}$ has enough projectives. Similarly, one proves that the category $\mathcal{O}_\downarrow$ has enough projectives (using Lemma 4.7). But then the equivalence of Lemma 4.7 implies that category ${}^1\mathcal{O}$ has enough injectives, as well. $\square$

Now, fix an associative algebra A with triangular structure $(A^\pm, \delta)$. Let ${}^1\mathcal{O}_\downarrow(A-A)$ be the category of finitely-generated left $A \otimes A^{\text{op}}$ -modules M such that the action on M of the subalgebras $A^- \otimes 1 \subset A \otimes A^{\text{op}}$ and $1 \otimes (A^+)^{\text{op}} \subset A \otimes A^{\text{op}}$ are both locally-nilpotent.

Lemma 4.12. *If conditions (2.6) (i.e., the assumptions of Duflo Theorem) hold then*

- (i) *The category ${}^1\mathcal{O}_\downarrow(A-A)$ is abelian and any object $M \in {}^1\mathcal{O}_\downarrow(A-A)$ has finite length.*
- (ii) *Any simple object of ${}^1\mathcal{O}_\downarrow(A-A)$ has the form $L = L' \boxtimes L''$, where L' is a simple object in ${}^1\mathcal{O}$ and L'' is a simple object in $\mathcal{O}_\downarrow$.*
- (iii) *The action of the element $\delta \otimes 1 + 1 \otimes \delta$ on any object $M \in {}^1\mathcal{O}_\downarrow(A-A)$ is locally-finite.*

³ this observation has been used earlier, see [Gu].

Proof. First, an argument very similar to the proof of Lemma 4.2 shows that analogues of statements (4.1)(1)-(3) hold for any object of ${}^1\mathcal{O}_\downarrow(A-A)$. Specifically, we leave to the reader to adapt the arguments that have been used above to prove that

- The action of the element $\delta \otimes 1 + 1 \otimes \delta$ on any object $M \in {}^1\mathcal{O}_\downarrow(A-A)$ is locally-finite.
- Each generalized eigenspace of the $(\delta \otimes 1 + 1 \otimes \delta)$ -action in $M \in {}^1\mathcal{O}_\downarrow(A-A)$ is finite dimensional, and the real parts of the generalized eigenvalues are all bounded from below.
- For any good filtration on an $A \otimes A^{\text{op}}$ -module $M \in {}^1\mathcal{O}_\downarrow(A-A)$, the module $\mathbf{gr} M$ is finitely-generated over $\mathbf{gr}(A^+) \otimes \mathbf{gr}(A^-)^{\text{op}}$.

Thus, part (iii) of the Lemma follows. We now prove part (ii). Fix a simple object $L \in {}^1\mathcal{O}_\downarrow(A-A)$. Choose a non-zero element $v \in L$ in the generalized δ -eigenspace corresponding to an eigenvalue with minimal possible real part. Commutation relations show that the action of any element of either $\mathfrak{n}^- \otimes 1$ or $1 \otimes \mathfrak{n}^+$ decreases the real part of the eigenvalue. Hence, we must have: $(\mathfrak{n}^- \otimes 1)v = 0$ and $(1 \otimes \mathfrak{n}^+)v = 0$. Therefore, the assignment $a'(v') \boxtimes a''(v'') \mapsto (a' \otimes a'')(v)$ gives a well-defined $A \otimes A^{\text{op}}$ -module map $\pi : L' \boxtimes L'' \rightarrow L$, where we put $L' := A/A \cdot \mathfrak{n}^-$ and $L'' := A^{\text{op}}/A^{\text{op}} \cdot \mathfrak{n}^+$, and where $v' \in L'$, $v'' \in L''$ denote the canonical generators (corresponding to the coset of $1 \in A$). The map π is nonzero, hence surjective, since L is simple.

Further, we know that $L' \in {}^1\mathcal{O}$ and $L'' \in \mathcal{O}_\downarrow$ (by Lemma 4.6 and its $\mathcal{O}_\downarrow$ -analogue). By our assumptions, all objects of ${}^1\mathcal{O}$ and $\mathcal{O}_\downarrow$ have finite length, cf. Corollary 4.8. Therefore both L' and L'' have finite Jordan-Hölder series with simple subquotients, say, $L'_1, \dots, L'_p$, and $L''_1, \dots, L''_q$, respectively. It is clear from Jacobson Density theorem that, for any $i \in [1, p]$, $j \in [1, q]$, the $A \otimes A^{\text{op}}$ -module $L'_i \boxtimes L''_j$ is simple. Hence, $L' \boxtimes L''$ also has a finite Jordan-Hölder series (as an $A \otimes A^{\text{op}}$ -module) with simple subquotients of the form $L'_i \boxtimes L''_j$. It follows that L , being a simple quotient of $L' \boxtimes L''$, must be isomorphic to some simple subquotient of the Jordan-Hölder series for $L' \boxtimes L''$, that is to some $L'_i \boxtimes L''_j$. Part (ii) follows.

We see from part (ii) that the category ${}^1\mathcal{O}_\downarrow(A-A)$ has finitely many isomorphism classes of simple objects. Then, mimicing the proof of Theorem 2.5(ii), one deduces that any object in ${}^1\mathcal{O}_\downarrow(A-A)$ has finite length. This completes the proof of part (i). $\square$

5 Jacquet functor and Proof of Theorem 2.3

We review the construction of Jacquet functor following [Ca] and [CO].

Let A be an associative algebra and $\mathfrak{r} \subset A$ a finite dimensional Lie subalgebra (with respect to the commutator bracket). Assume further that $\mathfrak{r}$ has the form of a semi-direct product $\mathfrak{n} \ltimes \mathbb{C} \cdot \delta$ such that

- (i) The adjoint action of $\mathfrak{n}$ on A is locally-nilpotent and

(ii) $\text{ad}\delta$ -action on A is locally-finite, and all eigenvalues of $\text{ad}\delta$ -action on $\mathfrak{n}$ are in $\mathbb{Z}_+$.

Note that the second condition implies that $\mathfrak{n}$ is a nilpotent Lie algebra. Let $\mathcal{U}_+$ denote the associative subalgebra (without unit) in A generated by $\mathfrak{n}$. Thus, $\mathcal{U}_+$ is a quotient of the augmentation ideal in the enveloping algebra $\mathcal{U}\mathfrak{n}$.

For each $k \geq 1$, let $\mathcal{U}_+^k$ be the k -th power of $\mathcal{U}_+$. These powers form a descending chain of subalgebras $\mathcal{U}_+ \supset \mathcal{U}_+^2 \supset \dots$, such that $\bigcap_{k \geq 1} \mathcal{U}_+^k = 0$. Note that condition (i) above implies that, for any $a \in A$, there exists $n = n(a) \in \mathbb{Z}$ such that

$$a \cdot \mathcal{U}_+^k \subset \mathcal{U}_+^{k+n} \cdot A \quad , \quad \forall k \gg 0. \quad (5.1)$$

Recall further that for any left A -module M , the linear dual $M^* = \text{Hom}_{\mathbb{C}}(M, \mathbb{C})$ has a natural right A -module structure. Given a *left* A -module M we define

$$\begin{aligned} \mathfrak{J}^\dagger(M) &:= \varprojlim_k (M/\mathcal{U}_+^k \cdot M)^* \\ &= \{m^* \in M^* \mid m^*(\mathcal{U}_+^k \cdot M) = 0 \text{ for some } k = k(m^*) \gg 0\}. \end{aligned} \quad (5.2)$$

Formula (5.1) insures that $\mathfrak{J}^\dagger(M)$ is an A -submodule of the right A -module M^* . Moreover, it is clear that the action on $\mathfrak{J}^\dagger(M)$ of the Lie subalgebra $\mathfrak{n} \subset A$ is locally nilpotent. Thus, we get a functor $\mathfrak{J}^\dagger : \text{left } A\text{-modules} \longrightarrow \text{right } A\text{-modules, locally-nilpotent relative to } \mathfrak{n}$.

It is convenient to get the following slightly different interpretation of this functor. Observe that formula (5.1) implies that the multiplication map: $A \xrightarrow{a \cdot} A$ (by any element $a \in A$) is continuous in the topology on A defined by the set $\{\mathcal{U}_+^k \cdot A\}_{k \geq 1}$ of fundamental neighborhoods of zero. Thus, the completion $\widehat{A} = \varprojlim_k A/\mathcal{U}_+^k \cdot A$ acquires the structure of

a complete topological algebra. There is a canonical algebra map: $A \rightarrow \widehat{A}$ with dense image. Similarly, for any left A -module M , the completion

$$\widehat{\mathfrak{J}}(M) = \varprojlim_k M/\mathcal{U}_+^k \cdot M \quad (5.3)$$

acquires the structure of a complete topological left $\widehat{A}$ -module. A non-commutative analogue of the standard Artin-Rees lemma implies that, on the category of finitely-generated A -modules, the functor $M \mapsto \widehat{\mathfrak{J}}(M)$ is exact. Furthermore, if M is finitely generated over A , then one has a canonical surjective morphism $\widehat{A} \otimes_A M \rightarrow \widehat{\mathfrak{J}}(M)$, see [AM, ch.10]. It is clear that $\mathfrak{J}^\dagger(M)$ is nothing but the *continuous dual* of $\widehat{\mathfrak{J}}(M)$.

The object $\widehat{\mathfrak{J}}(M)$ is too large to be finitely generated over A . Using the element $\delta \in A$, that has not played any role so far, one may do better.

Notation. Given $\lambda \in \mathbb{C}$ and a locally-finite δ -action on a vector space V , we write $[V]_\lambda$ for the generalized δ -eigenspace corresponding to the eigenvalue λ . We will often view A -modules as either $\mathcal{U}\mathfrak{r}$ - or $\mathcal{U}\mathfrak{n}$ -modules via the homomorphism $\mathcal{U}\mathfrak{r} \rightarrow A$ induced by the imbedding $\mathfrak{r} \hookrightarrow A$.

Let $\text{Mod}(A : \mathcal{U}_+)$ be the category of left A -modules which are finitely generated as $\mathcal{U}_+$ -modules. This is an abelian category since $\mathcal{U}_+$, being a quotient of the augmentation ideal in $\mathcal{U}\mathfrak{n}$, is Noetherian. Notice that for any $M \in \text{Mod}(A : \mathcal{U}_+)$ and any $k \geq 1$, the space

$M/\mathcal{U}_+^k \cdot M$ is finite dimensional. The action of $\delta \in A$ on M induces a δ -action on each finite dimensional space in the inverse system

$$M/\mathcal{U}_+ \cdot M \longleftarrow M/\mathcal{U}_+^2 \cdot M \longleftarrow M/\mathcal{U}_+^3 \cdot M \longleftarrow \dots \quad (5.4)$$

One deduces from the positivity of ad -eigenvalues on $\mathfrak{n}$ that, for each $\lambda \in \mathbb{C}$, the corresponding inverse system of generalized λ -eigenspaces: $[M/\mathcal{U}_+^k \cdot M]_\lambda$ stabilizes, i.e. the projections in (5.4) induce isomorphisms: $[M/\mathcal{U}_+^{k+1} \cdot M]_\lambda \xrightarrow{\sim} [M/\mathcal{U}_+^k \cdot M]_\lambda$, for all k sufficiently large.

Definition 5.5. Let $\mathfrak{J}(M) = \bigoplus_{\lambda \in \mathbb{C}} \left[\varprojlim_k M/\mathcal{U}_+^k \cdot M \right]_\lambda$ be the direct sum of all such “stable” generalized δ -eigenspaces, to be called the Jacquet module for M .

It is clear that $\mathfrak{J}(M)$ may be identified with the direct sum of all generalized δ -eigenspaces in $\widehat{\mathfrak{J}}(M)$. Thus, we have

- (1) $\mathfrak{J}(M)$ is an A -submodule in $\widehat{\mathfrak{J}}(M)$ which is dense in $\widehat{\mathfrak{J}}(M)$ in the $\mathcal{U}_+$ -adic topology.
- (2) δ -action on $\mathfrak{J}(M)$ is locally finite, with finite-dimensional generalized eigenspaces.
- (3) $\mathfrak{J}(M)$ is finitely-generated over $\mathcal{U}_+$, that is, $\mathfrak{J}(M) \in \text{Mod}(A : \mathcal{U}_+)$.
- (4) The set of δ -eigenvalues in $\mathfrak{J}(M)$ is bounded from below by some constant $\lambda(M) \in \mathbb{C}$.
- (5) $\mathfrak{J}^\dagger(M) = (\mathfrak{J}(M))^*$ is the restricted dual of $\mathfrak{J}(M)$.

Properties (1) and (2) above follow from the stabilization of eigen-spaces in the inverse system (5.4). To prove (3), let $N_0 \subsetneq N_1 \subsetneq \dots$ be a strictly ascending chain of $\mathcal{U}_+$ -submodules in $\mathfrak{J}(M)$. Property (2) implies that, for each $i \geq 0$, there exists $\lambda(i) \in \mathbb{C}$ such that $[N_i]_{\lambda(i)} \neq [N_{i+1}]_{\lambda(i)}$. It follows that for the corresponding closures in $\widehat{\mathfrak{J}}(M)$ we have: $\overline{N}_i \neq \overline{N}_{i+1}$. Thus, $\overline{N}_0 \subsetneq \overline{N}_1 \subsetneq \dots$ is a strictly ascending chain of $\widehat{\mathcal{U}}_+$ -submodules in $\widehat{\mathfrak{J}}(M)$. But this contradicts the fact that $\widehat{\mathfrak{J}}(M)$ is a Noetherian $\widehat{\mathcal{U}}_+$ -module (the latter holds since M is finitely-generated over $\mathcal{U}_+$, hence $\widehat{\mathfrak{J}}(M)$ is finitely-generated over $\widehat{\mathcal{U}}_+$). Thus, $\mathfrak{J}(M)$ must be a finitely-generated $\mathcal{U}_+$ -module, and (3) is proved. Property (4) follows from (3), and property (5) is clear from (2) and (4).

Since taking (generalized) eigenspaces is an exact functor, we conclude that the assignment $M \mapsto \mathfrak{J}(M)$ gives an exact functor $\mathfrak{J} : \text{Mod}(A : \mathcal{U}_+) \rightarrow \text{Mod}(A : \mathcal{U}_+)$, called the Jacquet functor.

Next, let $(A^\pm, \delta)$ be a triangular structure on A . We have the corresponding categories ${}^\dagger\mathcal{O}, \mathcal{O}_\dagger$. We also consider the category $\text{Mod}(A : \mathcal{U}_+)$ for the subalgebra $\mathcal{U}_+ := A^+$.

Lemma 5.6. (i) The Jacquet functor gives an exact functor $\mathfrak{J} : \text{Mod}(A : \mathcal{U}_+) \longrightarrow {}^\dagger\mathcal{O}$.

(ii) If conditions (2.6) hold, then the functor $\mathfrak{J}^\dagger$ gives an exact functor: $\text{Mod}(A : \mathcal{U}_+) \longrightarrow \mathcal{O}_\dagger$ and, moreover, $\mathfrak{J}^\dagger(M) = (\mathfrak{J}(M))^*$, for any $M \in \text{Mod}(A : \mathcal{U}_+)$.

Proof. To prove (i) we note that for any $M \in \text{Mod}(A : \mathcal{U}_+)$, the A^- -action on $\mathfrak{J}(M)$ is locally-nilpotent since the set of the real parts of δ -eigenvalues in $\mathfrak{J}(M)$ is bounded from

below by some constant $\lambda(M) \in \mathbb{R}$. We know also that $\mathfrak{J}(M)$ is finitely-generated over A^+ (property (3) above), hence, over A . Thus, $\mathfrak{J}(M) \in {}^1\mathcal{O}$, and (i) is proved.

Now, Lemma 4.7 yields $(\mathfrak{J}(M))^* \in \mathcal{O}_1$. But property (5) above says: $\mathfrak{J}^\dagger(M) \simeq (\mathfrak{J}(M))^*$. This completes the proof of part (ii). $\square$

Associated with the triangular structure $(A^\pm, \delta)$ on A , we have the subalgebra $A^+ \otimes (A^-)^{\text{op}} \subset A \otimes A^{\text{op}}$. We introduce an abelian category $\text{Mod}(A-A, A^+-A^-)$ of left $A \otimes A^{\text{op}}$ -modules as follows. If the triangular structure is *commutative*, we let $\text{Mod}(A-A, A^+-A^-)$ be the category of $A \otimes A^{\text{op}}$ -modules which are finitely generated over the subalgebra $A^+ \otimes (A^-)^{\text{op}}$. Thus, an object of $\text{Mod}(A-A, A^+-A^-)$ is an $(A-A)$ -bimodule, finitely-generated as an (A^+-A^-) -bimodule.

If the triangular structure on A is *non-commutative*, then the imbedding $A^+ \otimes (A^-)^{\text{op}} \hookrightarrow A \otimes A^{\text{op}}$ induces a graded algebra homomorphism: $\text{gr } A^+ \otimes \text{gr } A^- \hookrightarrow \text{gr } (A \otimes A^{\text{op}})$. We define $\text{Mod}(A-A, A^+-A^-)$ to be the category of finitely-generated left $A \otimes A^{\text{op}}$ -modules M such that $\text{gr } M$ is a finitely generated $(\text{gr } A^+) \otimes (\text{gr } A^-)$ -module, for any *good* increasing filtration on M . In particular, any object of $\text{Mod}(A-A, A^+-A^-)$ is a finitely-generated $A^+ \otimes (A^-)^{\text{op}}$ -module.

Recall that, given a non-commutative triangular structure on A , we have the Lie subalgebras $\mathfrak{n}^\pm \subset A^\pm$. In case of a *commutative* triangular structure on A , we let $\mathfrak{n}^\pm$ be any δ -stable finite-dimensional subspace of $A^\pm$ that generates $A^\pm$ as an associative algebra. We then regard $\mathfrak{n}^\pm$ as abelian Lie subalgebras in A . Thus, for any (commutative or non-commutative) triangular structure on A , we may consider the Lie algebra $\mathfrak{n}^+ \oplus \mathfrak{n}^- = \mathfrak{n}^+ \otimes 1 + 1 \otimes \mathfrak{n}^- \subset A \otimes A^{\text{op}}$. Therefore, there is a well-defined Jacquet functor on the category $\text{Mod}(A-A, A^+-A^-)$ (relative to the Lie algebra $\mathfrak{n}^+ \oplus \mathfrak{n}^- \subset A \otimes A^{\text{op}}$, and the element $\delta \otimes 1 + 1 \otimes \delta$). Furthermore, one mimics the proof of Lemma 5.6 to show an analogue of the Lemma, saying that the Jacquet functor is exact and takes the category $\text{Mod}(A-A, A^+-A^-)$ into ${}^1\mathcal{O}_1(A-A)$. Thus, we get an exact functor $\mathfrak{J} : \text{Mod}(A-A, A^+-A^-) \longrightarrow {}^1\mathcal{O}_1(A-A)$.

Proposition 5.7. (i) *The Jacquet functor $\mathfrak{J} : \text{Mod}(A-A, A^+-A^-) \longrightarrow {}^1\mathcal{O}_1(A-A)$ is faithful, that is we have: $M \in \text{Mod}(A-A, A^+-A^-)$ and $M \neq 0 \implies \mathfrak{J}(M) \neq 0$.*

(ii) *If conditions 2.6 hold for either ${}^1\mathcal{O}$ or $\mathcal{O}_1$, then any object of the category $\text{Mod}(A-A, A^+-A^-)$ has finite length.*

Remark. There is also a ‘one-sided’ analogue of this Proposition formulated as follows. Assume, for concreteness, that the triangular structure on A is non-commutative. We define $\text{Mod}(A, A^+)$ to be the category of finitely-generated left A -modules M such that $\text{gr } M$ is a finitely generated $(\text{gr } A^+)$ -module, for any *good* increasing filtration on M . In particular, any object of $\text{Mod}(A, A^+)$ is a finitely-generated A^+ -module, but the category $\text{Mod}(A, A^+)$ is, in general, *smaller* than the category $\text{Mod}(A : \mathcal{U}_+)$ for $\mathcal{U}_+ = A^+$. Similarly to Proposition 5.7 one proves

Proposition 5.8. (i) *The Jacquet functor gives a faithful functor $\mathfrak{J} : \text{Mod}(A, A^+) \rightarrow {}^1\mathcal{O}$.*

(ii) *If conditions 2.6 hold for ${}^1\mathcal{O}$ then any object of the category $\text{Mod}(A, A^+)$ has finite length.* $\square$

Our proof of Proposition 5.7 (and Proposition 5.8) will be based on Gabber's Separation theorem, which we now recall, cf. [Ga] for more details.

Let $\mathfrak{r} = \mathfrak{n} \rtimes \mathbb{C} \cdot \delta$ be a solvable Lie algebra such that the adjoint δ -action on $\mathfrak{n}$ is diagonal with all the eigenvalues in $\mathbb{Z}_+$, as at the beginning of this section. Write $\mathcal{U}_{\bullet} \mathfrak{r}$ and $\mathcal{U}_{\bullet} \mathfrak{n}$ for the standard increasing filtrations on the corresponding enveloping algebras. Thus, $\mathrm{gr}(\mathcal{U} \mathfrak{n}) = \mathrm{Sym} \mathfrak{n}$. Let M be a left $\mathcal{U} \mathfrak{r}$ -module. The following is ([Ga], Theorem 1):

Theorem 5.9 (Separation theorem). *Let M be a non-zero $\mathcal{U} \mathfrak{r}$ -module, and $\{M_j\}_{j \geq 0}$ an increasing filtration on M compatible with $\mathcal{U} \mathfrak{r}$ -action (i.e., $\mathcal{U}_i \mathfrak{r} \cdot M_j \subset M_{i+j}$, $\forall i, j \geq 0$). If $\mathrm{gr} M$ is finitely generated over $\mathrm{gr}(\mathcal{U} \mathfrak{n})$ (not only over $\mathrm{gr}(\mathcal{U} \mathfrak{r})$), then: $\mathfrak{n} \cdot M \neq M$.*

Remark. If $\mathfrak{n}$ is abelian, then the Separation theorem simplifies, and becomes the following standard result in Commutative Algebra (cf. e.g. [SW]): *If M is an $\mathcal{U} \mathfrak{r}$ -module which is finitely-generated over the subalgebra $\mathcal{U} \mathfrak{n}$, then: $\mathfrak{n} \cdot M \neq M$* (no filtration is needed in this case). To prove this, we claim first that the point $0 \in \mathrm{Spec}(\mathrm{Sym} \mathfrak{n})$ belongs to $\mathrm{Supp}(M)$. If not, then there is a polynomial $P \in \mathrm{Sym} \mathfrak{n}$ that vanishes on $\mathrm{Supp} M$ and such that $P(0) \neq 0$. Replacing P by its high enough power we may achieve that P annihilates M , i.e. $P \in \mathrm{Ann}(M) \subset \mathrm{Sym} \mathfrak{n}$. But the space $\mathrm{Ann}(M)$ is clearly stable under the adjoint δ -action on $\mathrm{Sym} \mathfrak{n}$. Moreover, since $P = P(0) + P_1$, where $P_1 \in \mathfrak{n} \cdot (\mathrm{Sym} \mathfrak{n})$, and all weights of $\mathrm{ad} \delta$ -action on $\mathfrak{n} \cdot \mathrm{Sym} \mathfrak{n}$ are strictly positive, we deduce from $P \in \mathrm{Ann}(M)$ that $P(0), P_1 \in \mathrm{Ann}(M)$. Since $P(0) \neq 0$ this yields $1 \in \mathrm{Ann}(M)$, a contradiction. Thus we have proved $0 \in \mathrm{Supp} M$. But then $M/\mathfrak{n} \cdot M$, the geometric fiber of M at 0, is non-zero, due to Nakayama lemma. Thus, $M \neq \mathfrak{n} \cdot M$. $\square$

Here is an example⁴ showing that the separation result may fail, in general, if (for $\mathfrak{n}$ non-commutative) the assumption of Theorem 5.9: ' $\mathrm{gr} M$ is finitely generated over $\mathrm{gr}(\mathcal{U} \mathfrak{n})$ ' is replaced by the weaker assumption: ' M is finitely generated over $\mathcal{U} \mathfrak{n}$ '.

Example. Let $\mathfrak{n}$ be the 3-dimensional Heisenberg Lie algebra with basis x, y, z , where z is central and $[x, y] = z$. Define a semi-direct product $\mathfrak{r} = \mathfrak{n} \rtimes \mathbb{C} \cdot \delta$ by the commutation relations: $[\delta, x] = x$, $[\delta, y] = y$, and $[\delta, z] = 2z$. Further, let D be the associative algebra of polynomial differential operators on the 2-plane $\mathbb{A} := \mathbb{C}^2$, with coordinates ' y, z '. The assignment:

$$x \mapsto z \frac{\partial}{\partial y}, \quad y \mapsto y, \quad z \mapsto z, \quad \delta \mapsto 2z \frac{\partial}{\partial z} + y \frac{\partial}{\partial y}.$$

extends to a Lie algebra imbedding: $\mathfrak{r} \hookrightarrow D$. We set $f(y, z) := y \cdot z - 1$. Thus $f \in \mathbb{C}[y, z]$ is a polynomial, and the equation $f = 0$ defines a hyperbola $X \subset \mathbb{A}$ such that

- The curve X is smooth, irreducible, and does not intersect the axis $z = 0$, and
- The restriction to X of the projection $\mathbb{A} \twoheadrightarrow \mathbb{C}$ along the y -axis gives an étale morphism: $X \rightarrow \mathbb{C}$.

Set $U := \mathbb{A} \setminus X$, an affine Zariski open subset in $\mathbb{C}^2$. The vector space $M := \mathbb{C}[U]/\mathbb{C}[\mathbb{A}] = \mathbb{C}[\mathbb{A}][\frac{1}{f}]/\mathbb{C}[\mathbb{A}]$ has a natural D -module structure (making M a simple holonomic D -module on the plane $\mathbb{A}$, with support X). It is straightforward to verify that the

⁴kindly communicated to me by O. Gabber

action of $\frac{\partial}{\partial y}$ on M induces isomorphisms:

$$\frac{1}{f^n} \cdot \mathbb{C}[y, z] \Big/ \frac{1}{f^{n-1}} \cdot \mathbb{C}[y, z] \xrightarrow{\sim} \frac{1}{f^{n+1}} \cdot \mathbb{C}[y, z] \Big/ \frac{1}{f^n} \cdot \mathbb{C}[y, z] \quad , \quad \forall n = 1, 2, \dots$$

It follows easily that $M = \mathcal{U}\mathfrak{n} \cdot [\frac{1}{f}]$, i.e., the class $[\frac{1}{f}] \in \mathbb{C}[\mathbb{A}][\frac{1}{f}]/\mathbb{C}[\mathbb{A}]$ generates M over the subalgebra $\mathcal{U}\mathfrak{n} \subset D$.

On the other hand, it is clear that $z \cdot M = M$, i.e., the separation property *fails* for M (in this case, $\mathbf{gr}(M)$ is not finitely generated over $\mathbf{gr}(\mathcal{U}\mathfrak{n})$). $\diamond$

Proof of Proposition 5.7. The argument is quite standard. Set $\mathfrak{n} = \mathfrak{n}^+ \oplus \mathfrak{n}^-$, a Lie subalgebra in $A \otimes A^{\text{op}}$, and write $\mathcal{U}_+$ for the augmentation ideal in $\mathcal{U}(\mathfrak{n}^+ \oplus \mathfrak{n}^-)$. The algebra $A^+ \otimes (A^-)^{\text{op}}$ is a quotient of $\mathcal{U}_+$. Hence we may (and will) view any $A \otimes A^{\text{op}}$ -module M as an $\mathcal{U}_+$ -module. We have a functor $\widehat{\mathfrak{J}}$ on $A \otimes A^{\text{op}}$ -modules defined as the completion with respect to the subalgebra $A^+ \otimes (A^-)^{\text{op}}$. In view of the identifications above, for any $M \in \text{Mod}(A-A, A^+-A^-)$, we can write: $\widehat{\mathfrak{J}}(M) = \varprojlim_k M/\mathcal{U}_+^k \cdot M$. The kernel

of the canonical map $M \rightarrow \widehat{\mathfrak{J}}(M)$ is clearly equal to $K = \bigcap_{k \geq 0} \mathcal{U}_+^k M$. It is routine to verify that K is an $A \otimes A^{\text{op}}$ -submodule in M . Moreover, a non-commutative version of Artin-Rees lemma, see [AM], [Ca], implies that $\mathcal{U}_+ \cdot K = K$ or, equivalently, that $\mathfrak{n} \cdot K = K$.

We would like to apply the Separation theorem to conclude that $K = 0$. To verify the conditions of that Theorem, observe that $K \in \text{Mod}(A-A, A^+-A^-)$, as a submodule of $M \in \text{Mod}(A-A, A^+-A^-)$. Choose a good filtration $K_\bullet$ on K . Then, since $\delta \in A_1$, for any $i \in \mathbb{Z}$, we get: $(\delta \otimes 1 + 1 \otimes \delta) \cdot K_i \subset K_{i+1}$. We see that all the conditions of the Separation theorem 5.9 hold. Hence $K = 0$, as we wanted.

Thus, the canonical map $M \rightarrow \widehat{\mathfrak{J}}(M)$ is injective. Hence, $M \in \text{Mod}(A-A, A^+-A^-)$ & $M \neq 0$ implies $\widehat{\mathfrak{J}}(M) \neq 0$. We conclude that $\mathfrak{J}(M)$, being a dense subspace in $\widehat{\mathfrak{J}}(M)$, is also non-zero, and part (i) follows.

To prove (ii), it suffices to verify the descending chain condition for any object $M \in \text{Mod}(A-A, A^+-A^-)$. Let $M = M^0 \supsetneq M^1 \supsetneq M^2 \supsetneq \dots$ be a descending chain of submodules in M . Using the exactness of Jacquet functor and part (i), from $M^i/M^{i+1} \neq 0$, $\forall i$, we deduce: $\mathfrak{J}(M^i)/\mathfrak{J}(M^{i+1}) = \mathfrak{J}(M^i/M^{i+1}) \neq 0$. Hence, $\mathfrak{J}(M^0) \supsetneq \mathfrak{J}(M^1) \supsetneq \mathfrak{J}(M^2) \supsetneq \dots$ is a strictly descending chain of submodules in $\mathfrak{J}(M)$. This contradicts the fact that $\mathfrak{J}(M)$, being an object of ${}^1\mathcal{O}_\downarrow(A-A)$, must have finite length by Lemma 4.12. Claim (ii) follows. $\square$

Proof of Theorem 2.4. Conditions $(T0)$, $(T0')$ imply that $A \in \text{Mod}(A-A, A^+-A^-)$, as an $(A-A)$ -bimodule. Thus, Theorem 2.4 follows directly from Proposition 5.7(ii). $\square$

Proof of Theorem 2.3. The implication (i) $\implies$ (ii) is standard.

To prove (ii) $\implies$ (iii), fix a prime ideal $I \subset A$. As we have observed above, $A \in \text{Mod}(A-A, A^+-A^-)$, hence $A/I \in \text{Mod}(A-A, A^+-A^-)$. Therefore, we may apply the Jacquet functor to get an object $\mathfrak{J}(A/I) \in {}^1\mathcal{O}_\downarrow(A-A)$.

Given an $A \otimes A^{\text{op}}$ -module, resp. $(A-A)$ -bimodule, M , we write $\mathbf{LAnn}(M)$ for the annihilator of M in the subalgebra $A \otimes 1 \subset A \otimes A^{\text{op}}$. The proof of Proposition 5.7 shows that A/I injects into $\widehat{\mathfrak{J}}(A/I)$ and, moreover, the image of A/I is dense in $\widehat{\mathfrak{J}}(A/I)$. Thus

we have $I = \mathbf{LAnn}(A/I) = \mathbf{LAnn}(\widehat{\mathfrak{J}}(A/I))$. Furthermore, $\mathbf{LAnn}(\widehat{\mathfrak{J}}(A/I)) = \mathbf{LAnn}(\mathfrak{J}(A/I))$, for $\mathfrak{J}(A/I)$ is dense in $\widehat{\mathfrak{J}}(A/I)$. Thus, $\mathbf{LAnn}(\mathfrak{J}(A/I)) = I$ is a prime ideal in A .⁵

Now, any object of ${}^1\mathcal{O}_\downarrow(A-A)$, in particular, $M = \mathfrak{J}(A/I)$, has finite length by Lemma 4.12. Let $0 = M_0 \subset M_1 \subset M_2 \subset \dots \subset M_n = M$, be its Jordan-Hölder series. For each $i = 1, 2, \dots, n$, put $J_i := \mathbf{LAnn}(M_i/M_{i-1})$. Then, by a standard argument, see e.g. [BG], one proves: $I = J_m$, for some $m \in [1, n]$.

We provide more details for the reader's convenience. First, since $J_i(M_i) \subset M_{i-1}$ we deduce:

$$J_1 \cdot \dots \cdot J_{n-1} \cdot J_n(M) \subset J_1 \cdot \dots \cdot J_{n-1}(M_{n-1}) \subset \dots \subset J_1(M_1) = 0.$$

Hence, the ideal $J_1 \cdot \dots \cdot J_n$ annihilates M , and therefore $J_1 \cdot \dots \cdot J_n \subset \mathbf{LAnn}(M)$. But, $\mathbf{LAnn}(M) = I$ being prime, the inclusion above implies that there exists an m such that $J_m \subset I$. On the other hand, since M_i/M_{i-1} is a subquotient of M , for any $i = 1, 2, \dots, n$, one clearly has an opposite inclusion $I = \mathbf{LAnn}(M) \subset \mathbf{LAnn}(M_i/M_{i-1}) = J_i$. Thus, we obtain $I = J_m = \mathbf{LAnn}(M_m/M_{m-1})$, as claimed.

By construction, M_m/M_{m-1} is a simple object of ${}^1\mathcal{O}_\downarrow(A-A)$. Lemma 4.12 yields $M_m/M_{m-1} = L' \boxtimes L''$, where $L' \in {}^1\mathcal{O}$ and $L'' \in \mathcal{O}_\downarrow$ are some simple objects. Further, it is clear that $\mathbf{LAnn}(L' \boxtimes L'') = \mathbf{LAnn}(L')$. Thus, we deduce that $I = \mathbf{LAnn}(M_m/M_{m-1}) = \mathbf{LAnn}(L' \boxtimes L'') = \mathbf{LAnn}(L')$ is the annihilator of a simple object of ${}^1\mathcal{O}$. Part (iii) follows.

Part (iv) is equivalent to (iii) by duality. The implication (iii) $\implies$ (i) is trivial. $\square$

6 Applications to Symplectic reflection algebras

Let (V, ω) be a finite dimensional symplectic vector space, and $\Gamma \subset Sp(V, \omega)$ a finite subgroup of symplectic automorphisms of V . An element $s \in \Gamma$ is called a *symplectic reflection* if $\text{rk}(\text{Id} - s) = 2$. Let S denote the set of symplectic reflections in Γ . The group Γ acts on S by conjugation. For each $s \in S$, there is an ω -orthogonal direct sum decomposition $V = \text{Image}(\text{Id} - s) \oplus \text{Ker}(\text{Id} - s)$, and we write ω_s for the (possibly degenerate) skew-symmetric form on V which coincides with ω on $\text{Image}(\text{Id} - s)$, and has $\text{Ker}(\text{Id} - s)$ as the radical.

Write $\mathbb{C}\Gamma$ for the group algebra of Γ , and $(TV) \# \Gamma$ for the smash-product of the tensor algebra TV with $\mathbb{C}\Gamma$. From now on we identify V with V^* via the symplectic form, hence the Symmetric algebra on V gets identified with $\mathbb{C}[V]$, the polynomial algebra on V .

In [EG], we have introduced a class of associative algebras $H_{t,c}(V, \omega, \Gamma)$, called Symplectic reflection algebras, as follows. Fix an $\text{Ad } \Gamma$ -invariant function $c : S \rightarrow \mathbb{C}$, $s \mapsto c_s$, and a complex number $t \in \mathbb{C}$. Let $\kappa : V \times V \rightarrow \mathbb{C}\Gamma$ be a skew-symmetric $\mathbb{C}$ -bilinear pairing given by the formula:

$$\kappa(x, y) = t \cdot \omega(x, y) \cdot 1 + \sum_{s \in S} c_s \cdot \omega_s(x, y) \cdot s \quad , \quad \forall x, y \in V.$$

⁵It was pointed out by a referee that a similar observation is contained in [J2, n.2.4].

We define the Symplectic reflection algebra with parameters (t, c) by

$$\mathbf{H}_{t,c}(V, \omega, \Gamma) := (TV \# \Gamma) / I \langle x \otimes y - y \otimes x - \kappa(x, y) \in T^2V \oplus \mathbb{C}\Gamma \rangle_{x,y \in V}, \quad (6.1)$$

where $I \langle \dots \rangle$ stands for the two-sided ideal in $(TV) \# \Gamma$ generated by the indicated set. Thus, $\mathbf{H}_{t,c}(V, \omega, \Gamma)$ is an associative algebra which may be thought of as a deformation of $\mathbb{C}[V] \# \Gamma$ (= cross-product of $\mathbb{C}[V]$ with Γ). Clearly, $\mathbf{H}_{t,c}(V, \omega, \Gamma)$ contains $\mathbb{C}\Gamma$ as a subalgebra. Furthermore, there is a natural increasing filtration on $\mathbf{H}_{t,c}(V, \omega, \Gamma)$ such that $\mathbb{C}\Gamma$ has filtration degree zero, and elements of $V \subset TV$ are assigned filtration degree one. It has been proved in [EG] that the corresponding associated graded algebra is described by the following canonical graded algebra isomorphism:

$$\text{gr } \mathbf{H}_{t,c}(V, \omega, \Gamma) \simeq \mathbb{C}[V] \# \Gamma \quad (\text{Poincaré-Birkhoff-Witt property}). \quad (6.2)$$

Write $\mathbf{e} = \frac{1}{\#\Gamma} \sum_{g \in \Gamma} g \in \mathbb{C}\Gamma$ for the averaging idempotent, viewed as an element in $\mathbf{H}_{t,c}(V, \omega, \Gamma)$. We let $\mathbf{eHe} := \mathbf{e} \cdot \mathbf{H}_{t,c}(V, \omega, \Gamma) \cdot \mathbf{e}$ be the *spherical subalgebra* in $\mathbf{H}_{t,c}(V, \omega, \Gamma)$. The increasing filtration on $\mathbf{H}_{t,c}(V, \omega, \Gamma)$ induces a filtration on the spherical subalgebra, and from (6.2) one finds that $\text{gr}(\mathbf{eHe}) = \mathbb{C}[V]^\Gamma = \mathbb{C}[V/\Gamma]$, the algebra of Γ -invariant polynomials on V . The canonical Poisson structure on the commutative algebra $\text{gr}(\mathbf{eHe})$ makes V/Γ a Poisson variety. It is easy to see, cf. ([EG], Theorem 1.6), that this Poisson structure on V/Γ equals the t -multiple of the standard one induced by the symplectic structure on V (for $t = 0$ the spherical subalgebra $\mathbf{eHe}$ becomes commutative, and the corresponding Poisson structure on $\text{gr}(\mathbf{eHe}) = \mathbb{C}[V/\Gamma]$ reduces to zero).

Symplectic leaves in V/Γ . The leaves of the standard Poisson structure on $\mathbb{C}[V/\Gamma]$ are described as follows, see [BrGo]. Let $\text{Isotropy}(\Gamma, V)$ denote the (finite) set of all subgroups $G \subset \Gamma$ that occur as isotropy groups of points in V . Given a subgroup $G \subset \Gamma$, write $V^G \subset V$ for the vector subspace of G -fixed points. Let V_G denote the image of V^G under the projection $V \twoheadrightarrow V/\Gamma$, and let $\mathring{V}_G = V_G \setminus \bigcup'_{G' \in \text{Isotropy}(\Gamma, V)} (V_G \cap V_{G'})$, be the complement in V_G of the union of all *proper* subsets in V_G of the form $V_G \cap V_{G'}$ (here the symbol $\bigcup'$ indicates that the union is taken over proper subsets only). It is easy to see that for each subgroup $G \in \text{Isotropy}(\Gamma, V)$, the set $\mathring{V}_G$ equals the image of the subset $\mathring{V}^G \subset V^G$ formed by the points whose isotropy group equals G . If N_G denotes the normalizer of G in Γ , then the group N_G/G acts freely on $\mathring{V}^G$ and we have $\mathring{V}_G \simeq \mathring{V}^G / (N_G/G)$. It follows that each set $\mathring{V}_G$ is a smooth connected locally-closed subvariety of V/Γ and one shows that these varieties are exactly the symplectic leaves in V/Γ , see e.g., [BrGo]. Hence, there are only finitely many symplectic leaves.

From the Irreducibility theorem 2.1 we deduce

Corollary 6.3. *Let $\mathbf{H}_{t,c}(V, \omega, \Gamma)$ be a Symplectic reflection algebra, and $t \neq 0$. Then, for any primitive ideal I in the spherical subalgebra $\mathbf{e} \cdot \mathbf{H}_{t,c}(V, \omega, \Gamma) \cdot \mathbf{e}$, the variety $\mathcal{V}(I)$ has the form V_G , for a certain subgroup $G \in \text{Isotropy}(\Gamma, V)$.*

We remark that Theorem 2.1 is not applicable to the Symplectic reflection algebra $\mathbf{H}_{t,c}(V, \omega, \Gamma)$ itself since $\text{gr } \mathbf{H}_{t,c}(V, \omega, \Gamma) = \mathbb{C}[V] \# \Gamma$, is a non-commutative algebra. Note

however, that finitely generated $\mathbb{C}[V]\#\Gamma$ -modules may be naturally identified with Γ -equivariant coherent sheaves on V . The support of such a sheaf is a Γ -stable subvariety in V . In particular, given a two-sided ideal $I \subset \mathbf{H}_{t,c}(V, \omega, \Gamma)$ one can view $\mathbf{gr}(\mathbf{H}_{t,c}(V, \omega, \Gamma)/I)$ as a finitely generated $\mathbb{C}[V]\#\Gamma$ -module, via (6.2). Let $\mathcal{V}(I) \subset V$ stand for its support.

Using the technique of *Poisson orders* developed in [BrGo], one can refine our argument to obtain the following

Proposition 6.4. *For any primitive ideal $I \subset \mathbf{H}_{t,c}(V, \omega, \Gamma)$, the variety $\mathcal{V}(I)$ is the Γ -saturation of vector subspace V^G , for a certain subgroup $G \in \text{Isotropy}(\Gamma, V)$. $\square$*

Rational Cherednik algebras. Let W be a finite Coxeter group in a complex vector space $\mathfrak{h}$. Thus, $\mathfrak{h}$ is the complexification of a real Euclidean vector space and W is generated by a finite set $S \subset W$ of reflections $s \in S$ with respect to certain hyperplanes $\{H_s\}_{s \in S}$ in that Euclidean space.

For each $s \in S$, we choose a nonzero linear function $\alpha_s \in \mathfrak{h}^*$ which vanishes on H_s (called the *positive root* corresponding to s), and let $\alpha_s^\vee = 2(\alpha_s, -)/(\alpha_s, \alpha_s) \in \mathfrak{h}$ be the corresponding coroot. The group W acts on the set S , and diagonally on $V = \mathfrak{h} \oplus \mathfrak{h}^*$, by conjugation. We equip the space $V = \mathfrak{h} \oplus \mathfrak{h}^*$ with the canonical symplectic structure of the cotangent bundle.

Following [EG], to each W -invariant function $c : S \rightarrow \mathbb{C}$, $c \mapsto c_s$, one associates the Symplectic reflection algebra $\mathbf{H}_c := \mathbf{H}_{1,c}(V, \omega, W)$, called the *rational Cherednik algebra*. The algebra $\mathbf{H}_c$ is generated by the vector spaces $\mathfrak{h}$, $\mathfrak{h}^*$, and the set W , with defining relations (cf. formula (1.15) of [EG] for $t = 1$) given by

$$\begin{aligned} w \cdot x \cdot w^{-1} &= w(x) \quad , \quad w \cdot y \cdot w^{-1} = w(y) \quad , \quad \forall y \in \mathfrak{h} \quad , \quad x \in \mathfrak{h}^* \quad , \quad w \in W \\ x_1 \cdot x_2 &= x_2 \cdot x_1 \quad , \quad y_1 \cdot y_2 = y_2 \cdot y_1 \quad , \quad \forall y_1, y_2 \in \mathfrak{h} \quad , \quad x_1 \quad , \quad x_2 \in \mathfrak{h}^* \\ y \cdot x - x \cdot y &= \langle y, x \rangle + \sum_{s \in S} c_s \cdot \langle y, \alpha_s \rangle \langle \alpha_s^\vee, x \rangle \cdot s \quad , \quad \forall y \in \mathfrak{h} \quad , \quad x \in \mathfrak{h}^* . \end{aligned} \tag{6.5}$$

Thus, the elements $x \in \mathfrak{h}^*$ generate a subalgebra $\mathbb{C}[\mathfrak{h}] \subset \mathbf{H}_c$ of polynomial functions on $\mathfrak{h}$, the elements $y \in \mathfrak{h}$ generate a subalgebra $\mathbb{C}[\mathfrak{h}^*] \subset \mathbf{H}_c$, and the elements $w \in W$ span a copy of the group algebra $\mathbb{C}W$ sitting naturally inside $\mathbf{H}_c$. Furthermore, the Poincaré-Birkhoff-Witt property yields a ‘triangular decomposition’ $\mathbf{H}_c \simeq \mathbb{C}[\mathfrak{h}^*] \otimes \mathbb{C}[W] \otimes \mathbb{C}[\mathfrak{h}]$, see ([EG], Corollary 4.4). We put $A^+ := \mathbb{C}^{>0}[\mathfrak{h}]^W$, and $A^- := \mathbb{C}^{>0}[\mathfrak{h}^*]^W$. Further, in [BEG] we have constructed an element $\delta \in \mathbf{H}_c$ such that the data $(A^\pm, \delta)$ gives a commutative triangular structure on the rational Cherednik algebra $\mathbf{H}_c$.

The categories ${}^1\mathcal{O}, \mathcal{O}_1$ (see Definition 2.2) for $\mathbf{H}_c$ have only finitely many (isomorphism classes of) simple objects, by [DO]. Thus Theorem 2.3 applies (note that we are using here only the easy, purely commutative, part of Gabber’s Separation theorem, since $A^\pm$ are commutative algebras) and we get

Corollary 6.6. *Both Duflo theorem and Theorem 2.4 hold for rational Cherednik algebras.*

In [BEG] we introduced a category $\mathcal{O}(\mathbf{H}_c)$ to be the category of finitely-generated left $\mathbf{H}_c$ -modules M such that the A^- -action on M is locally-finite. It is clear from the

triangular decomposition $H_c \simeq \mathbb{C}[\mathfrak{h}^*] \otimes \mathbb{C}[W] \otimes \mathbb{C}[\mathfrak{h}]$ that in the notation of §4 we have: ${}^1\mathcal{O} \subset \mathcal{O}(H_c) \subset \text{Mod}(H_c, A^+)$. Note that the inclusion ${}^1\mathcal{O} \subset \mathcal{O}(H_c)$ is *strict* since we do not require the locally-finite A^- -action to be locally-nilpotent. We have also introduced in [BEG] a category $\mathcal{HC}$ of Harish-Chandra bimodules over the Cherednik algebra H_c , and showed that (in our present notation) any object of $\mathcal{HC}$ belongs to $\text{Mod}(A-A, A^+-A^-)$. Thus, from Propositions 5.7-5.8 we deduce

Corollary 6.7. *Any object of the categories $\mathcal{O}(H_c)$ and $\mathcal{HC}$ has finite length.* $\square$

Remark. Notice that in [BEG] we were only able to prove that any object of the category ${}^1\mathcal{O}$ has finite length, but the technique in loc. cit. was insufficient to prove Corollary 6.7 in full generality.

As has been explained before Proposition 6.4, for any H_c -module, hence for an object $M \in \mathcal{O}(H_c)$, one can define a W -stable subvariety $\text{Supp } M \subset \mathfrak{h} \oplus \mathfrak{h}^*$. The A^- -action on M being locally-finite, it follows, since $A^- = \mathbb{C}^{>0}[\mathfrak{h}^*]^W$, that $\text{Supp } M \subset \mathfrak{h} \oplus \{0\}$. Next, identify $\mathfrak{h}^*$ with $\mathfrak{h}$ via the invariant form and observe that, since W is finite, the isotropy group of a point $(y, x) \in \mathfrak{h} \oplus \mathfrak{h}^* = \mathfrak{h} \oplus \mathfrak{h}$ coincides with the isotropy group of the generic linear combination $t_1 \cdot y + t_2 \cdot x$, $t_1, t_2 \in \mathbb{C}$. Therefore we see that, in the notation of Theorem 6.3, the set $\text{Isotropy}(W, \mathfrak{h} \oplus \mathfrak{h}^*)$ is exactly the set $\text{Parab}(W)$ of all parabolic subgroups in W , cf. [Hu].

Theorem 6.8. *For any simple object $M \in \mathcal{O}(H_c)$, the variety $\text{Supp } M$ is the W -saturation of vector subspace $\mathfrak{h}^G \oplus \{0\}$, for a certain subgroup $G \in \text{Parab}(W)$.*

Sketch of Proof. We adapt the known argument used in the case of highest weight modules over $\mathcal{U}\mathfrak{g}$, the enveloping algebra of a semisimple Lie algebra. To this end, write $\text{Ann } M \subset H_c$ for the annihilator of $M \in \mathcal{O}(H_c)$. Thus, $\text{Ann } M$ is a primitive ideal in H_c .

Introduce the notation $\Lambda := \text{Supp } M \subset \mathfrak{h} \oplus \mathfrak{h}^*$. Clearly, $\Lambda \subset \text{Supp}(H_c/\text{Ann } M)$. Further, by Proposition 6.4 there exists a subgroup $G \in \text{Parab}(W)$ such that $\text{Supp}(H_c/\text{Ann } M) = W \cdot (\mathfrak{h} \oplus \mathfrak{h}^*)^G$. Therefore we get:

$$\Lambda \subset (\mathfrak{h} \oplus \{0\}) \cap W \cdot (\mathfrak{h} \oplus \mathfrak{h}^*)^G = W \cdot (\mathfrak{h}^G \oplus \{0\}) \quad (6.9)$$

Next, we claim that

$$\dim \Lambda = \dim(\mathfrak{h}^G). \quad (6.10)$$

To prove this, choose a finite-dimensional $\mathbb{C}[\mathfrak{h}^*]^W$ -stable subspace $M^0 \subset M$ that generates M over the subalgebra $\mathbb{C}[\mathfrak{h}]^W$. For each $i \geq 0$, set $M_i = (H_c)_i \cdot M^0$, where $\{(H_c)_i\}_{i \geq 0}$ is the standard increasing filtration on the Cherednik algebra H_c , cf. [EG]. This gives a filtration on M compatible with that on H_c and such that $\text{gr } M$ is finitely-generated over the subalgebra $\text{gr } \mathbb{C}[\mathfrak{h}]^W \subset \text{gr } H_c$. Hence, the restriction to $\text{Supp } M \subset \text{Spec}(\text{gr } H_c)$ of the canonical projection: $\text{Spec}(\text{gr } H_c) \rightarrow \text{Spec}(\mathbb{C}[\mathfrak{h}]^W)$ is a finite morphism. Therefore the dimension of the variety $\text{Supp } M$ is unaffected by replacing the algebra H_c by the algebra $\mathbb{C}[\mathfrak{h}]^W$. Now, following Joseph [J3], we define an increasing chain of $\mathbb{C}[\mathfrak{h}]^W$ -submodules $D_k(M) \subset \text{End}_{\mathbb{C}} M$, $k = 0, 1, \dots$, inductively as follows

$$D_0(M) = \mathbb{C}[\mathfrak{h}]^W, \quad \text{and} \quad D_k(M) = \{u \in \text{End}_{\mathbb{C}} M \mid [u, \mathbb{C}[\mathfrak{h}]^W] \subset D_{k-1}(M)\}.$$

Set $D(M) := \bigcup_{k \geq 0} D_k(M)$. Joseph showed that $D(M)$ is an associative subalgebra in $\mathbf{End}_{\mathbb{C}} M$ (which should be thought of as an algebra of ‘differential operators’ on M). Moreover, he proved in ([J3], Lemma 2.3) that $\dim \mathcal{S}upp D(M) \leq 2 \dim(\mathcal{S}upp M)$.

Observe further that the action in M of an element $a \in \mathbf{H}_c$ gives an endomorphism $\hat{a} \in \mathbf{End}_{\mathbb{C}} M$, and the assignment $a \mapsto \hat{a}$ gives an algebra imbedding: $\mathbf{H}_c / \mathbf{Ann} M \hookrightarrow \mathbf{End}_{\mathbb{C}} M$. The adjoint action of the subalgebra $\mathbb{C}[\mathfrak{h}]^W$ on $\mathbf{H}_c$ being locally-nilpotent, see [BEG], the image of the embedding: $\mathbf{H}_c / \mathbf{Ann} M \hookrightarrow \mathbf{End}_{\mathbb{C}} M$ is contained in $D(M)$. From this, following Joseph [J3], one derives:

$$\dim \mathcal{S}upp(\mathbf{H}_c / \mathbf{Ann} M) \leq \dim \mathcal{S}upp D(M) \leq 2 \dim(\mathcal{S}upp M). \quad (6.11)$$

Hence, using (6.9) and writing ‘ $\dim(-)$ ’ instead of ‘ $\dim \mathcal{S}upp(-)$ ’, we deduce

$$\dim(\mathfrak{h} \oplus \mathfrak{h}^*)^G = \dim(\mathbf{H}_c / \mathbf{Ann} M) \leq 2 \dim M = 2 \dim \Lambda \leq 2 \dim \mathfrak{h}^G = \dim(\mathfrak{h} \oplus \mathfrak{h}^*)^G.$$

It follows that all the inequalities above must be equalities, and (6.10) is proved. In particular, there exists $\Lambda^\circ \subset (\mathfrak{h}^G \oplus \{0\})$, an irreducible component of Λ such that $\dim \Lambda^\circ = \dim(\mathfrak{h}^G \oplus \{0\})$. This dimension equality yields: $\Lambda^\circ = \mathfrak{h}^G \oplus \{0\}$.

We claim next that *all* irreducible components of $\Lambda = \mathcal{S}upp M$ have the same dimension, i.e., the following version of Gabber’s equi-dimensionality theorem holds for the Cherednik algebra

Proposition 6.12. *For any simple $\mathbf{H}_c$ -module M , the variety $\mathcal{S}upp M$ is equidimensional.*

To prove this, we recall that the equi-dimensionality theorem is, as explained e.g. in [Gi], a formal consequence of Gabber-Kashiwara theorem. The proof of the latter theorem given, e.g. in ([Gi], p.342-345) works for any filtered algebra $\mathcal{A}$ such that $\mathbf{gr} \mathcal{A}$ is the coordinate ring of a *smooth* affine algebraic variety. Now, in our present situation, we have: $\mathbf{gr} \mathbf{H}_c = \mathbb{C}[\mathfrak{h} \oplus \mathfrak{h}^*] \# W$, which is not a commutative algebra. However, the formal microlocalization construction can be carried out with respect to any multiplicative set $S \subset \mathcal{A}$, ($0 \notin S$), for any algebra $\mathcal{A}$, provided the principal symbols of the elements of S belong to the center of $\mathbf{gr} \mathcal{A}$, see ([Gi], footnote on p. 337). In our case we have a central subalgebra: $\mathbb{C}[\mathfrak{h} \oplus \mathfrak{h}^*]^W \subset \mathbf{gr} \mathbf{H}_c$. Furthermore, the algebra $\mathbf{gr} \mathbf{H}_c$ has *finite* homological dimension, as a cross-product of the polynomial algebra $\mathbb{C}[\mathfrak{h} \oplus \mathfrak{h}^*]$ with a finite group. Going through the proof of Gabber-Kashiwara theorem given in ([Gi], pp. 342–345) one sees that the finiteness of homological dimension of $\mathbf{H}_c$, plus the existence of the formal microlocalization with respect to elements whose symbols belong to $\mathbb{C}[\mathfrak{h} \oplus \mathfrak{h}^*]^W \setminus \{0\}$, is enough to conclude that the proof of Gabber-Kashiwara theorem applies to the algebra $\mathbf{H}_c$ as well. This implies Proposition 6.12.

From Proposition 6.12 we deduce that all irreducible components of the variety $\Lambda = \mathcal{S}upp M$ have the same dimension, which is equal to $\dim \mathfrak{h}^G$. But then the inclusion in (6.9) forces each irreducible component to be a W -translate of $\mathfrak{h}^G$, and Theorem 6.8 follows. $\square$

Remark. It was pointed out to us by A. Yekutieli, that Gabber’s equi-dimensionality theorem also holds for the spherical subalgebra $\mathbf{eHe}$ in any Symplectic reflection algebra $\mathbf{H} = \mathbf{H}_{t,c}(V, \omega, \Gamma)$, i.e. one has

Proposition 6.13. *Let $H_{t,c}(V, \omega, \Gamma)$ be a Symplectic reflection algebra. Then, for any simple $e\text{He}$ -module M , the variety $\text{Supp } M$ is equidimensional.*

Sketch of proof. Recall that PBW-property (6.2) implies that $\text{gr}(e\text{He}) = \mathbb{C}[V]^\Gamma = \mathbb{C}[V/\Gamma]$. We see that $\text{Spec}(\text{gr}(e\text{He})) \simeq V/\Gamma$ is not a smooth variety. Nonetheless, V/Γ is a Gorenstein variety, and homological duality formalism is known to work for Gorenstein varieties as nicely as for smooth varieties. In particular, it follows from [YZ] that the bimodule $e\text{He}$ is (up to shift) a rigid dualizing complex for the algebra $e\text{He}$ (in the sense of [YZ]). Therefore, the proof of Gabber-Kashiwara theorem still goes through, cf. [YZ, Purity theorem]. $\square$

References

- [AM] M. Atiyah, I. Macdonald: *Introduction to commutative algebra*. Addison-Wesley, Reading, Ma. 1969.
- [BEG] Yu. Berest, P. Etingof, and V. Ginzburg: *Cherednik algebras and differential operators on quasi-invariants*. [arXiv:math.QA/0111005](#), to appear in *Duke Math. J.*
- [Be] J. Bernstein: *Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients*. (Russian) *Funkcional. Anal. Appl.* **5** (1971), 1–16.
- [BG] J. Bernstein, S. Gelfand: *Tensor products of finite- and infinite-dimensional representations of semisimple Lie algebras*. *Compositio Math.* **41** (1980), 245–285.
- [BoBr] W. Borho, J.-L. Brylinski: *Differential operators on homogeneous spaces. I. Irreducibility of the associated variety for annihilators of induced modules*. *Invent. Math.* **69** (1982), 437–476.
- [BoKr] W. Borho, H. Kraft: *Über die Gelfand-Kirillov-Dimension*. *Math. Ann.* **220** (1976), 1–24.
- [BrGo] K. Brown, I. Gordon: *Poisson orders, symplectic reflection algebras and representation theory*. [arXiv:math.RT/020104](#).
- [Ca] W. Casselman: *Jacquet modules for real reductive groups*. *Proceedings of the ICM (Helsinki, 1978)*, pp. 557–563, *Acad. Sci. Fennica, Helsinki*, 1980.
- [CO] W. Casselman, M. Osborne: *The restriction of admissible representations to n* . *Mathem. Ann.* **233** (1978), 193–198.
- [Ch] I. Cherednik: *Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald operators*, *IMRN (Duke math.J.)* **9** (1992), 171–180.
- [CG] N. Chriss, V. Ginzburg: *Representation theory and complex geometry*. Birkhäuser Boston, 1997.
- [Di] J. Dixmier: *Enveloping algebras*. Graduate Studies in Math., **11** A.M.S., Providence, RI, 1996.
- [Du] M. Duflo: *Sur la classification des idéaux primitifs dans l’algèbre enveloppante d’une algèbre de Lie semi-simple*. *Ann. of Math.* **105** (1977), 107–120.
- [DO] C. Dunkl, E. Opdam: *Dunkl operators for complex reflection groups*. [arXiv:math.RT/0108185](#).
- [EG] P. Etingof, V. Ginzburg: *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*. *Invent. Math.* **147** (2002), 243–348, [[arXiv:math.AG/0011114](#)].
- [Ga] O. Gabber: *A separation theorem and Harish-Chandra modules*. Unpublished manuscript 1981.
- [Gal] O. Gabber: *The integrability of the characteristic variety*. *Amer. J. Math.* **103** (1981), 445–468.
- [Gi] V. Ginzburg: *Characteristic varieties and vanishing cycles*. *Invent. Math.* **84** (1986), 327–402.
- [Gu] N. Guay: *Projective modules in the category $\mathcal{O}$ for the Cherednik algebra*. Preprint, University of Chicago, December 2001.
- [Hu] J. Humphreys: *Reflection groups and Coxeter groups*. Cambridge Studies in Advanced Mathematics, **29** Cambridge University Press, Cambridge, 1990.

- [J1] A. Joseph: *Primitive ideals in enveloping algebras*. Proceedings of the ICM (Warsaw, 1983), 403–414, PWN, Warsaw, 1984.
- [J2] A. Joseph: *On the associated variety of a primitive ideal*. J. Algebra **93** (1985), 509–523.
- [J3] A. Joseph: *Gelfand-Kirillov dimension for the annihilators of simple quotients of Verma modules*. J. London Math. Soc. **18** (1978), 50–60.
- [KT] M. Kashiwara, T. Tanisaki: *The characteristic cycles of holonomic systems on a flag manifold related to the Weyl group algebra*. Invent. Math. **77** (1984), 185–198.
- [Po] A. Polishchuk: *Algebraic geometry of Poisson brackets*. J. Math. Sci. **84** (1997), 1413–1444.
- [SW] J.T. Stafford, N.R. Wallach: *The restriction of admissible modules to parabolic subalgebras*. Trans. Amer. Math. Soc. **272** (1982), 333–350.
- [Vo] D. Vogan: *Associated varieties and unipotent representations. Harmonic analysis on reductive groups*. 315–388, Progr. Math., **101**, Birkhäuser Boston, Boston, MA, 1991.
- [YZ] A. Yekutieli, J. Zhang: *Rings with Auslander dualizing complexes*. J. Algebra **213** (1999), 1–51.

Department of Mathematics, University of Chicago, Chicago IL 60637, USA;
 ginzburg@math.uchicago.edu